

Antioxidant enzyme induction in pea plants under high irradiance

J.A. HERNÁNDEZ*¹, C. ESCOBAR**, G. CREISSEN*** and P.M. MULLINEAUX****

*Dept. de Mejora Vegetal, Centro de Edafología y Biología Aplicada del Segura, CSIC, Apartado 164, E-30100 Espinardo-Murcia, Spain**

*Facultad de Ciencias del Medio Ambiente, Universidad de Castilla-La Mancha, E-45071 Toledo, Spain***

*Dept. Disease and Stress Biology, John Innes Centre, Colney, Norwich NR4 7UH, UK****

*Dept. Biological Science, University of Essex, Colchester, Essex CO4 3SQ, UK*****

Abstract

Exposure of pea plants to high irradiance (HI) for 60 min caused a reversible photoinhibition as shown by changes in variable to maximum fluorescence ratio (F_v/F_m). A significant decline in F_m was observed in leaves from both pea cultivars subjected to HI, the decrease being higher in JI281 than in JI399 plants. The values recovered during the post-stress period in both cultivars. In both cultivars, minimal fluorescence (F_0) increased under HI, but in cultivar JI399 F_0 recovered to initial value during the post-stress period. The expression of antioxidant enzyme genes was higher in JI399 than in JI281, both in control and stressed plants. In JI281, after 60 min of HI, an induction of the transcripts of *CAT*, *chlMDHAR*, *cytAPX* and *cytCu,Zn-SOD* was observed, whereas there was a slight increase in *PHGPX*, *stAPX* and *chlCu,Zn-SOD* mRNAs. After 24 h of the recovery period, the induction of some transcripts was not maintained (*CAT*, *cytAPX* and *cytCu,Zn-SOD*), whereas the induction of others was maintained (*PHGPX* and *chlCu,ZnSOD*) or even increased (*cytGR*, *stAPX* and *chlMDHAR*). In JI399, *CAT* and *cytAPX* were increased strongly after 60 min of HI, and slight increases were observed in *cytGR*, *chlGR* and *chlMDHAR*. In the post-stress period the expression of *stAPX*, *cytGR* and *chlMDHAR* was even slightly higher than after 60 min of HI, however, expression of *CAT*, *cytAPX*, *cytCu,ZnSOD*, *chlCu,ZnSOD* and *chlGR* decreased.

Additional key words: ascorbate peroxidase, catalase, gene expression, photooxidative stress, *Pisum sativum*, superoxide dismutase.

Introduction

High irradiance (HI) exposure is one of the most common causes of oxidative stress in plants (Dat *et al.* 2000). Under excess of irradiation, enzymatic processes for CO₂ fixation become rate-limiting and, as a result, photosynthesis produces more NADPH and ATP than necessary. This accumulation of redox and energy equivalents will decrease the plastoquinone pool and/or inhibit the water-splitting complex, inevitably leading to PS 2 inactivation, the so-called photoinhibition (Anderson *et al.* 1997, Karpinski *et al.* 1997, 1999).

There are reports on the changes in activity and expression of antioxidant enzymes in response to HI stress (Gillham and Dodge 1987, Foyer *et al.* 1989, Karpinski *et al.* 1997, Yoshimura *et al.* 2000, Hernández *et al.* 2004) but the results vary according to plant materials and treatment conditions employed.

The protection of plants against AOS produced in excess during adverse environmental conditions is achieved by means of different strategies and, in particular, by partial suppression of its production and scavenging of the AOS already produced (Murgia *et al.* 2004). During HI stress, free radicals are formed in PS 2, which are harmful to the photosynthetic apparatus (Aro *et al.* 1993). Partial suppression of AOS production in the chloroplasts during HI stress is achieved by degradation of the D1 protein of PS 2 leading to its inactivation (Anderson *et al.* 1997, Karpinski *et al.* 1997, 1999).

Plants contain a complex antioxidant system to detoxify AOS that includes carotenoids, ascorbate, glutathione, tocopherols, anthocyanin pigments and enzymes such as superoxide dismutase, catalase, glutathione peroxidase, peroxidases and the enzymes

Received 14 July 2004, accepted 20 May 2005.

Abbreviations: AOS - activated oxygen species; APX - ascorbate peroxidase; ASC-GSH cycle - ascorbate-glutathione cycle; CAT - catalase; CuZn-SOD - copper,zinc-containing superoxide dismutase; DHAR - dehydroascorbate reductase; F_0 , F_m , F_v - parameters of chlorophyll fluorescence; HI - high irradiance; GR - glutathione reductase; GPX - glutathione peroxidase; H₂O₂ - hydrogen peroxide; MDHAR - monodehydroascorbate reductase; SOD - superoxide dismutase; O₂^{•-} - superoxide radical; PHGPX - phospholipid hydroperoxide glutathione peroxidase.

¹ Corresponding author; fax: (+34) 968 396213, e-mail: jahernan@cebas.csic.es

involved in the ascorbate-glutathione cycle (Foyer and Halliwell 1976): ascorbate peroxidase, dehydroascorbate reductase, monodehydroascorbate reductase and glutathione reductase.

Materials and methods

Plants, growth and treatments: *Pisum sativum* L. (cvs. JI281 and JI399) seedlings, individually planted in pots, were grown in a controlled-environment growth chamber [18-h photoperiod, irradiance of $200 \pm 25 \mu\text{mol}(\text{photon}) \text{m}^{-2} \text{s}^{-1}$, temperature of $15 \pm 1.5^\circ\text{C}$ and relative humidity of $75 \pm 5\%$]. JI281 is a semi-domesticated land race of pea from Ethiopia whereas JI399 is a typical domesticated garden pea cultivar (Ellis *et al.* 1992).

Experiments were performed with 21-d-old pea plants exposed to HI for up to 60 min ($4000 \mu\text{mol m}^{-2} \text{s}^{-1}$). To eliminate heat effects and to disperse light evenly, light was reflected by a mirror and directed through a frosted-glass filter filled with cold water. Control plants were exposed to $200 \pm 25 \mu\text{mol m}^{-2} \text{s}^{-1}$. After this period, some light-stressed plants were returned for 24 h to normal growth conditions (re-exposed to $200 \pm 25 \mu\text{mol m}^{-2} \text{s}^{-1}$) before sampling (poststress period). All leaves were frozen in liquid N_2 and stored at -70°C until use.

Measurement of chlorophyll fluorescence: The F_0 (minimal chlorophyll fluorescence), F_m (maximal chlorophyll fluorescence) and photosystem 2 efficiency parameter F_v/F_m (the ratio of variable to maximal fluorescence; $F_v = F_m - F_0$) were measured using a portable *Plant Efficiency Analyser* (PEA; *Hansatech Instrument*, King's Lynn, Norfolk, UK), on 10 min dark-adapted leaves at 90 % light saturation, according to the manufacturers' instructions.

Northern blotting: Total RNA from leaves was extracted as previously described (Creissen and Mullineaux 1995). Poly A^+ -mRNA was purified by chromatography, using oligo d(T) cellulose spin columns (*Amersham Biosciences*, Buckinghamshire, UK) according to the manufacturer's instructions. Poly A^+ -mRNA (3 - 5 μg)

In this work, we investigate the effect of HI on PS 2 efficiency in pea leaves, as well as on the levels of mRNAs encoding crucial enzymes of AOS metabolism.

was denatured, separated electrophoretically and transferred onto a nitrocellulose membrane as described in Hernández *et al.* (2000). The loading of an equivalent amount of Poly A^+ -mRNA for each time point was checked on gels stained with ethidium bromide and they showed equal intensities (data not shown). Northern blots were visualised by autoradiography after hybridisation with ^{32}P -labelled DNA probes. Hybridisation was carried out in 0.3 M sodium phosphate buffer pH 7.2, containing 1 mM EDTA, 7 % SDS and 1 % BSA, at 65°C (homologous probes) or 55°C (heterologous probes). Washing was with $0.1\times$ SSC (65°C) for homologous probes and with $1\times$ SSC or $2\times$ SSC for heterologous probes. Northern blots were performed twice and representative blots are shown.

The specific probes used were: cytosolic and chloroplastic Cu,Zn-SOD from *Nicotiana plumbaginifolia* (Bowler *et al.* 1989, Van Camp *et al.* 1990), cytosolic cytGR (*GOR 2*, Stevens *et al.* 1997) and chloroplastic GR (*GOR 1*, Creissen and Mullineaux 1995), catalase from tobacco (Willekens *et al.* 1994), *PHGPX* (Mullineaux *et al.* 1998), chlMDAR from *Arabidopsis* (Genbank accession number T04550), cytosolic APX from pea (Santos *et al.* 1996) and stromal APX from spinach (Ishikawa *et al.* 1995).

The filters were exposed to X-ray film and/or were visualised on a *BAS 1000 Phosphorimager* analyser (*Fuji Photofilm Co.*, Kanagawa, Japan). Scanning values were calculated by *BASIS* software (*Fuji Photofilm Co.*) installed on the *BAS 1000*.

Statistics: Comparisons among means were made using Duncan's multiple range test, calculated at $P < 0.05$. Statistical procedures were carried out with the software package *SPSS 11.0* for Windows.

Results

Exposure of 3-week-old pea plants at high irradiance ($4000 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 60 min caused a significant photoinhibition in both cultivars as indicated by the decline in the PS 2 efficiency F_v/F_m (Fig. 1A). After 24 h at $200 \mu\text{mol m}^{-2} \text{s}^{-1}$, the efficiency of PS 2 recovered as shown by the increase in F_v/F_m (Fig. 1A) to values of the control plants. In both cultivars, HI caused an increase in F_0 . In JI281, F_0 partially recovered after 24 h, but still was increased in relation to time zero. However, in JI399, F_0 values reached control values after 24 h (Fig. 1B). A

significant decline in F_m was observed in leaves from both cultivars subjected to HI, the decrease being greater in JI281 than in JI399 plants. F_m values recovered during the post-stress period in both cultivars.

The expression of antioxidant enzyme genes was higher in cultivar JI399 than in JI281, both in control and in stressed plants (Fig. 2). In cultivar JI281, after 1 h of HI, an induction of the transcripts of *CAT*, *chlMDHAR*, *cytAPX* and *cytCu,Zn-SOD* was observed, whereas there were a slight increases in *PHGPX*, *stAPX* and




Fig 1. Effect of HI on F_v/F_m ratios (A), F_0 (B) and F_m (C). The F_v/F_m ratio, F_0 and F_m were measured in three individual plants obtained from three independent experiments ($n = 9$). The seedlings were grown at irradiance $200 \mu\text{mol m}^{-2} \text{s}^{-1}$. In the middle of the photoperiod, 3-week-old plants were exposed to HI ($4000 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 60 min and returned to low irradiance. Means with different letters are significantly different according to Duncan's multiple range test ($P = 0.05$).

Table 1. Transcript levels relative to the time 0 of genes coding antioxidant enzymes from attached pea leaves exposed to HI. A representative Northern blot is shown. Values were obtained after the scanning of RNA gel blot hybridization with cDNA probes. Value at time 0 is set to 1.

Probe	JI-281		JI-399	
	HI	Post-stress	HI	Post-stress
chlMDHAR	2.04	3.33	1.40	1.64
GOR 2	1.35	2.22	1.21	1.50
GOR 1	0.96	0.81	1.20	0.70
CAT1	2.00	1.30	2.24	1.15
PHGPX	1.71	1.93	1.04	0.80
cytAPX	4.16	1.4	2.03	0.73
stAPX	1.4	3.15	0.50	1.40
chlCuZnSOD	1.53	1.33	0.97	0.60
cytCuZnSOD	3.36	1.95	0.80	0.62

Discussion

Short-term exposure of pea plants to HI caused a significant photoinhibition, although F_v/F_m values had recovered after 24 h of the post-stress period. A similar response in photosynthesis has been described in spinach




Fig 2. Northern blot hybridisation analysis of poly (A+) RNA from pea plants exposed to HI. Names of probes used are indicated on the left. A representative Northern blot is shown.

chlCu,Zn-SOD mRNAs. After 24 h of the recovery, some transcripts were not maintained (*CAT*, *cytAPX* and *cytCuZn-SOD*), whereas others were maintained (*PHGPX* and *chlCu,ZnSOD*) or even increased (*cytGR*, *stAPX* and *chlMDHAR*) (Fig. 2, Table 1).

In cultivar JI399, *CAT* and *cytAPX* were increased after 1 h of HI, and slight increases were observed in *cytGR*, *chlGR* and *chlMDHAR* (Fig. 2, Table 1). After 24 h recovery, the expression of *CAT*, *stAPX*, *cytGR* and *chlMDHAR* was slightly higher than after 60 min of HI (Fig. 2, Table 1). However, contents of *cytAPX*, *cytCu,ZnSOD*, *chlCu,ZnSOD* and *chlGR* transcripts were lower when compared to those exhibited in control plants (time zero) and after 1 h of HI stress (Fig. 2, Table 1).

and *Arabidopsis* plants as well as in detached pea leaves subjected to HI stress (Yoshimura *et al.* 2000, Karpinski *et al.* 1997, Hernández *et al.* 2004). The increase in F_0 during HI has been described also in *Arabidopsis* plants

as well as in detached pea leaves (Karpinski *et al.* 1997, Hernández *et al.* 2004). Elevated F_0 has been considered as reflecting thylakoid membrane disturbance and photoinhibitory damage, and decreases in F_v/F_m and F_m during HI indicate that photoinhibition of photosynthesis occurred (Krause and Weiss 1991, Balachandran and Osmond 1994, Karpinski *et al.* 1997, Bertamini and Nedunchezian 2004).

mRNAs from some genes encoding antioxidant enzymes accumulated in both cultivars during HI, although to different extents. At time zero, as well as in stressed and recovered plants, expression of all mRNAs was higher in cultivar JI399 than in JI281. This could contribute to a more efficient response to the AOS that may be generated in chloroplasts and cytosol during the HI and post-stress periods. However, after HI, the increases observed for *chlMDHAR*, *cytCuZnSOD* and *cytAPX* were higher in JI281 than in JI399 plants, which could be due to the lower AOS generation in JI399 compared to JI281 after HI, as shown previously in detached leaves from the same cultivars (Hernández *et al.* 2004).

The strong induction of *CAT*, *cytAPX* and *cytCu,ZnSOD* in JI281 indicates that HI could have increased O_2^- and H_2O_2 levels in the cytosol. In the same way, the induction of *CAT* and *cytAPX* in JI399 could indicate also an increased H_2O_2 generation in the cytosolic compartment. An increase in H_2O_2 has been observed also in detached leaves from these pea cultivars after 60 min of HI, being higher in JI281 than in JI399 (Hernández *et al.* 2004).

On the other hand, the higher expression of *chlMDHAR*, *chlGR*, *chlPHGPX*, *stAPX* and *chlCu,Zn-SOD* observed in JI399 under HI, could suggest a higher protection against AOS in chloroplasts from this pea cultivar. Probably, the greater alterations in F_0 and F_m observed in JI281 could be related with the lower expression of all antioxidant enzymes analysed.

In detached pea leaves and in *Arabidopsis* plants, photoinhibition was correlated with an increase in H_2O_2 (Karpinski *et al.* 1997, Hernández *et al.* 2004). A similar response has been observed in spinach, where a transient increase in H_2O_2 levels was observed after 1 h of HI, correlated with a drop in F_v/F_m values (Yoshimura *et al.* 2000). So, the decrease in F_v/F_m and F_m , as well as the increase in F_0 , observed in pea plants subjected to HI could also be due to an increase in AOS in their chloroplasts. In bean plants, continuous irradiation induced premature senescence caused by enhanced production of AOS (Procházková and Wilhelmová 2004).

The increase in *cytGR* transcripts observed in pea leaves after 24 h post-stress has been previously

described in *Arabidopsis* plants and in detached pea leaves (Karpinski *et al.* 1997; Hernández *et al.* 2004), and a similar increase in *chlMDHAR* has been described also in detached pea leaves recovered from HI (Hernández *et al.* 2004). In wheat, DHAR activity declined during the early stages of seedling growth under a high-light regime and this implies that regeneration of ASC for scavenging of H_2O_2 is catalysed mostly by MDHAR and not by DHAR (Mishra *et al.* 1995). In the present study, we have not measured DHAR activity or expression, but the induction of *chlMDHAR* and the slight decrease in *chlGR*, in both pea cultivars, suggest that in chloroplasts, ASC seems to be regenerated mostly *via* MDHAR. Conversely, the increase in *cytGR* in recovered plants indicates that ASC could be regenerated mainly *via* GSH in the cytosolic compartment.

It has been suggested that in the induction of transcripts encoding for antioxidant enzymes under HI, both H_2O_2 and the redox status of ascorbate could be involved (Hernández *et al.* 2004). However, in *Arabidopsis* plants, a role for the redox state of glutathione and the plastoquinone pool also has been proposed (Karpinski *et al.* 1997). Therefore, in the present study, and although no measurements of H_2O_2 , ascorbate or glutathione levels have been carried out, the induction of transcripts encoding for antioxidant enzymes in response to HI could also have been mediated by AOS or by the alteration in the redox state of ascorbate and/or glutathione.

The data obtained about the expression of genes encoding antioxidant enzymes suggest that HI could alter AOS levels, both in the chloroplasts and cytosol of pea leaves. The induction of cytosolic antioxidant enzymes could provide an alternative protection when the chloroplastic defence systems are compromised under light stress conditions (Karpinski *et al.* 1997, Hernández *et al.* 2004). Support for this was reported by Yoshimura *et al.* (2000) who found that stress conditions, including HI, drought, salinity and methyl viologen treatments, which enhanced AOS in chloroplasts, also induces the cytosolic scavenging system.

In conclusion, results indicated that HI caused a reversible photoinhibition of photosynthesis in pea chloroplasts, and that changes in both F_0 and F_m contributed to the decrease in F_v/F_m , the changes being more important in JI281 than in JI399 leaves. Also, it seems that HI produces an increase in AOS that could regulate the accumulation of mRNAs encoding antioxidant enzymes. These data also suggest that pea cultivar JI399 seems to be relatively more tolerant than JI281 to HI, at least partly due to its higher expression levels of antioxidant enzymes.

References

- Anderson, J.M., Park, Y.I., Chow, W.S.: Photoinactivation and photoprotection of photosystem II in nature. - *Physiol. Plant.* **100**: 214-223, 1997.
- Aro, E.M., McCaffery, S., Anderson, J.M.: Photoinhibition and D1 protein degradation in peas acclimated to different growth irradiances. - *Plant Physiol.* **103**: 835-843, 1993.

- Balachandran, S., Osmond, C.B.: Susceptibility of tobacco leaves to photoinhibition following infection with two strains of tobacco mosaic virus under different light and nitrogen nutrition regimes. - *Plant Physiol.* **104**: 1051-1057, 1994.
- Bertamini, M., Nedunchezian, N.: Photosynthesis responses for *Vitis vinifera* plants grown at different photon flux densities under field conditions. - *Biol. Plant.* **48**: 149-152, 2004.
- Bowler, C., Alliotte, T., De Loose, M., Van Montagu, M., Inzé, D.: The induction of manganese superoxide dismutase in response to stress in *Nicotiana plumbaginifolia*. - *EMBO J.* **8**: 31-38, 1989.
- Creissen, G.P., Mullineaux, P.M.: Cloning and characterisation of glutathione reductase cDNAs and identification of two genes encoding the tobacco enzyme. - *Planta* **197**: 422-425, 1995.
- Dat, J., Van den Abeele, S., Vranová, E., Van Montagu, M., Inzé, D., Van Breusegem, F.: Dual action of the active oxygen species during plant stress responses. - *Cell. mol. Life Sci.* **57**: 779-795, 2000.
- Ellis, T.H.N., Turner, L., Hellens, R.P., Lee, D., Harker, C.L., Enard, C., Domoney, C., Davies, D.R.: Linkage maps in pea. - *Genetics* **130**: 649-663, 1992.
- Foyer, C.H., Dujardin, M., Lemoine, Y.: Responses of photosynthesis and the xanthophyll and ascorbate cycles to changes in irradiances, photoinhibition and recovery. - *Plant Physiol. Biochem.* **27**: 751-760, 1989.
- Foyer, C.H., Halliwell, B.: The presence of glutathione and glutathione reductase in chloroplasts: a proposed role in ascorbic acid metabolism. - *Planta* **133**: 21-25, 1976.
- Gillham, D.J., Dodge, A.D. Chloroplasts superoxide and hydrogen peroxide scavenging systems from pea leaves: seasonal variations. - *Plant Sci.* **50**: 105-109, 1987.
- Hernández, J.A., Jiménez, A., Mullineaux, P.M., Sevilla, F.: Tolerance of pea (*Pisum sativum* L.) to long-term salt stress is associated with induction of antioxidant defences. - *Plant Cell Environ.* **23**: 853-862, 2000.
- Hernández, J.A., Escobar, C., Creissen, G., Mullineaux, P.M.: Role of hydrogen peroxide and the redox state of ascorbate in the induction of antioxidant enzymes in pea leaves under excess light stress. - *Funct. Plant Biol.* **31**: 359-368, 2004.
- Ishikawa, T., Sakai, K., Takeda, T., Shigeoka, S.: Cloning and expression of cDNA encoding a new type of ascorbate peroxidase from spinach. - *FEBS Lett.* **367**: 28-32, 1995.
- Karpinski, S., Escobar, C., Karpinska, B., Creissen, G., Mullineaux, P.M.: Photosynthetic electron transport regulates the expression of cytosolic ascorbate peroxidase genes in *Arabidopsis* during excess light stress. - *Plant Cell* **9**: 627-640, 1997.
- Karpinski, S., Reynolds, H., Karpinska, B., Wingsle, G., Creissen, G., Mullineaux, P.M.: Systemic signaling and acclimation in response to excess excitation energy in *Arabidopsis*. - *Science* **284**: 654-657, 1999.
- Krause, G.H., Weiss, E.: Chlorophyll fluorescence and photosynthesis. - *Annu. Rev. Plant Physiol. Plant mol. Biol.* **42**: 313-349, 1991.
- Mishra, H.P., Fatma, T., Singhal, G.S.: Development of antioxidative defense system of wheat seedlings in response to high light. - *Physiol. Plant.* **95**: 77-82, 1995.
- Mullineaux, P.M., Karpinski, S., Jiménez, A., Cleary, S., Robinson, C., Creissen, G.: Identification of cDNAs encoding plastid-targeted glutathione peroxidase. - *Plant J.* **13**: 375-379, 1998.
- Murgia, I., Tarantino, D., Vannini, C., Bracale, M., Carravieri, S., Soave, C.: *Arabidopsis thaliana* plants overexpressing thylakoidal ascorbate peroxidase show increased resistance to paraquat-induced photooxidative stress and to nitric oxide-induced cell death. - *Plant J.* **38**: 940-953, 2004.
- Procházková, D., Wilhelmová, N.: Changes in antioxidant protection in bean cotyledons during natural and continuous irradiation-accelerated senescence. - *Biol. Plant.* **48**: 33-39, 2004.
- Santos, M., Gousseau, H., Lister, C., Foyer, C., Creissen, G., Mullineaux, P.M.: Cytosolic ascorbate peroxidase from *Arabidopsis thaliana* L. is encoded by a small multigene family. - *Planta* **198**: 64-69, 1996.
- Stevens, R.G., Creissen, G.P., Mullineaux, P.M.: Cloning and characterization of a cytosolic glutathione reductase cDNA from pea (*Pisum sativum* L.) and its expression in response to stress. - *Plant mol. Biol.* **35**: 641-654, 1997.
- Van Camp, W., Bowler, C., Villarroel, R., Tsang, E.W.T., Van Montagu, M., Inzé, D.: Characterization of iron superoxide dismutase cDNAs from plants obtained by genetic complementation in *Escherichia coli*. - *Proc. nat. Acad. Sci. USA* **87**: 9903-9907, 1990.
- Willekens, H., Langebartels, C., Tiré, C., Van Montagu, M., Inzé, D., Van Camp, W.: Differential expression of catalase genes in *Nicotiana plumbaginifolia* (L.). - *Proc. nat. Acad. Sci. USA* **91**: 10450-10454, 1994.
- Yoshimura, K., Yabuta, Y., Ishikawa, T., Shigeoka, S.: Expression of spinach ascorbate peroxidase isoenzymes in response to oxidative stress. - *Plant Physiol.* **123**: 223-233, 2000.