

The antioxidative role of anthocyanins in *Arabidopsis* under high-irradiance

Q. ZHANG¹, L.-J. SU¹, J.-W. CHEN², X.-Q. ZENG¹, B.-Y. SUN¹, and C.-L. PENG^{1*}

College of Life Science, South China Normal University, Guangzhou 510631, Guangdong, P.R. China¹

College of Agronomy and Biotechnology, Yunnan Agricultural University, Kunming 650201, Yunnan, P.R. China²

Abstract

To uncover the potential antioxidative role of anthocyanins *in vivo* in protecting photosynthetic tissues from photoinhibition, the effects of high irradiance [HI, 1300 $\mu\text{mol}(\text{photon})\text{m}^{-2}\text{s}^{-1}$] were studied using detached leaves derived from *Arabidopsis* wild-type (WT) and the mutant deficient in anthocyanin biosynthesis (*tt3tt4*). HI stress caused decreased chlorophyll content and photochemical efficiency, but increased cell-membrane leakage and contents of hydrogen peroxide and superoxide radical in the leaves of both *Arabidopsis* phenotypes, but the WT plants showed better HI tolerance than *tt3tt4* mutant. HI caused a significant increase in the 1,1-diphenyl-2-picrylhydrazyl scavenging capacity in WT but not in the *tt3tt4* mutant. The anthocyanins could not contribute substantially to light-shielding during the periods of HI stress, because the anthocyanin content in WT was very low and the colour of leaves was the same as in the *tt3tt4* mutant. Thus, it was assumed that the better HI tolerance in WT was mostly related to the potential antioxidative role of anthocyanins.

Additional key words: chlorophyll fluorescence imaging, hydrogen peroxide, membrane leakage, mutant, superoxide radical.

Introduction

Anthocyanins are plant pigments produced by the flavonoid biosynthetic pathway. They are often present in flowers, fruits, leaves and stems, ranging in colour from orange/red to purple/blue. Anthocyanins function in flowers and fruits primarily to attract pollinators and seed distributors (Chalker-Scott 1999, Gould and Lister 2006).

Many environmental factors (e.g. light, temperature, nutrition, drought and infection) have an effect on the synthesis of anthocyanins (De Jong 1991, Beggs and Wellmann 1994, Meng *et al.* 2004, Piovan and Filippini 2007, Ismail and Mohamed 2010). In recent years, further research in the functional roles of anthocyanins became possible with new research methods (Agati *et al.* 2007).

Recently, the role of anthocyanins in photoprotection has been proposed (Gould *et al.* 1995, Field *et al.* 2001, Hoch *et al.* 2001). According to this hypothesis, the photoprotective roles of anthocyanins can be fulfilled in two ways: either by simply screening visible radiation or/and by radical scavenging (Wang *et al.* 1997, Archetti

et al. 2009). The photoprotective roles, in which anthocyanins act as sunscreens to attenuate the visible radiation penetrating the mesophyll and thus reduce the excitation pressure, have been extensively reported (Smillie and Hetherington 1999, Field *et al.* 2001, Steyn *et al.* 2002, Manetas *et al.* 2003, Merzlyak *et al.* 2008). However, anthocyanins may also indirectly protect plants against excess radiation by their oxy-radical scavenging properties (Gould *et al.* 2002a, Steyn *et al.* 2002). Our previous study also indicated that leaves containing anthocyanins had a significantly greater antioxidant potential than did green leaves (Peng *et al.* 2006). Although some studies have confirmed that anthocyanins are powerful antioxidants (Tsuda *et al.* 1996, Wang *et al.* 1997, Gould *et al.* 2002a, Neill and Gould 2003), direct experimental evidence for the *in vivo* engagement of anthocyanins in antioxidative defense of leaves is lacking, and indirect indications are sparse (Manetas 2006).

Excess irradiance may be harmful for plants that are

Received 8 June 2010, accepted 16 March 2011.

Abbreviations: OH \cdot - hydroxyl radical; CHS - chalcone synthase; DAB - diaminobenzidine; DFR - dihydroflavono-4-reductase; DPPH - 1,1-diphenyl-2-picrylhydrazyl; ETR - apparent electron transport rate; F_v/F_m - variable to maximum fluorescence ratio (the maximum photosystem 2 quantum yield); NBT - nitroblue tetrazolium; O $_2^{\cdot-}$ - superoxide radical; qP - the coefficient of photochemical quenching; ROS - reactive oxygen species; *tt3tt4* - *tt3tt4* mutant deficient in anthocyanin biosynthesis; WT - wild-type; Φ_{PS2} - effective photosystem 2 quantum yield.

Acknowledgements: We thank Prof. Wah Soon Chow from the Australian National University for his useful suggestions on the manuscript. This research is financially supported by National Natural Science Foundation of China (30770173, 30870385), The State Key Basic Research and Development Plan of China (973 Program; 2009CB118504). The first two authors contributed equally to this work

* Corresponding author: fax: (+86-20) 85212078, e-mail: pengchl@scib.ac.cn

unable to balance the absorbed/utilized energy ratio (Huner *et al.* 1998). When the absorption of light energy exceeds the capacity of photosynthesis and the photoprotective mechanisms are overwhelmed, reactive oxygen species (ROS), such as superoxide radicals ($O_2^{\cdot-}$), hydrogen peroxide (H_2O_2) and hydroxyl radicals (OH^{\cdot}), can be produced (Havaux and Niyogi 1999), thereby resulting in oxidative stress. Hence, high irradiance can be

used as a specific generator of ROS. In the present study, the wild-type and the anthocyanin-deficient mutant of *Arabidopsis* (*tt3tt4*) were used to investigate the antioxidative potential of anthocyanins. Our working hypothesis was that anthocyanins *in vivo* function to relieve stress caused by high irradiance through their powerful antioxidative capacity.

Materials and methods

Seeds of wild-type (WT) *Arabidopsis thaliana* ecotype Landsberg erecta (*Ler*) and the Landsberg erecta mutant deficient in anthocyanin biosynthesis (*tt3tt4*), which is deficient in both the dihydroflavonol 4-reductase (DFR) locus and the chalcone synthase (CHS) locus (Shirley *et al.* 1995, Peer *et al.* 2001), were obtained from the *Arabidopsis* Biological Resource Center, Columbus, OH, USA. The seed coat of the WT is brown, while the coat of the mutants is yellow. Seeds were imbibed for 2 d at 4 °C in the dark to synchronise germination, and then sown on sterilised compost. Plant seedlings were grown routinely in a growth cabinet at temperature of 20 - 22 °C, 16-h photoperiod, irradiance of 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and relative humidity of 80 %. The third and fourth mature leaves from 25-d-old plants were used for the present study.

One set of leaves cut from the *tt3tt4* mutant and WT plants were floated in a pure-water bath at 20 °C, and the light sources were halogen lamps placed above a circulating-water tank with thermal insulation. The leaves were placed under high irradiance (HI, 1 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$) and were periodically taken for particular analyses.

After centrifugation of the methanolic extracts for 2 min at 3 000 g, the supernatant absorbance ($A_{530} - A_{657}$) was quantified by UV- visible spectrophotometer Lambda 24, Perkin-Elmer, Waltham, MA, USA) according to the methods described by Wade *et al.* (2003). In addition, absorption spectra were also recorded from 400 to 700 nm. Flavonoid and total phenolics extracted from leaf pieces were analyzed and estimated following the methods of Fukumoto and Mazza (2000). Chlorophyll (Chl) content was determined in 95 % ethanol extract as described by Lichtenthaler and Wellburn (1983).

The scavenging of 1,1-diphenyl-2-picrylhydrazyl (DPPH) was determined as described by Peng *et al.* (2000). In brief, 50 % ethanol extracts were prepared from the leaves, followed by centrifugation at 5 000 g for 15 min. The resulting supernatant was mixed with DPPH solution. The decrease in absorbance was measured at 525 nm using a spectrophotometer Lambda 25 (Perkin-Elmer).

For determination of cell-membrane leakage rate, leaves were immersed in double-distilled water for 1.5 h at room temperature, followed by a 30-min boiling. The conductivity of a solution of leaked electrolytes before and after boiling was determined using a DDS-11 A conductometer (Shanghai Dapu Instruments, Shanghai, China).

ROS localization was conducted according to

Romero-Puertas *et al.* (2004) and Zeng *et al.* (2010) with some modifications. After HI treatment for 0, 120 and 240 min, samples were collected for analysis of H_2O_2 and $O_2^{\cdot-}$ localization *in situ*. H_2O_2 was visualized by diaminobenzidine (DAB) staining. Samples from each treatment were immersed in a solution of DAB (1 mg cm^{-3} , Sigma-Aldrich, St. Louis, MI, USA) in 50 mM phosphate buffer (pH 7.0), vacuum-infiltrated for 10 min and then incubated at room temperature for 8 h in the dark. Then white light (80 $\mu\text{mol m}^{-2} \text{s}^{-1}$) was switched on until the appearance of brown spots, characteristic of the reaction product of DAB with H_2O_2 . Chlorophyll was then bleached by immersion in boiling ethanol (75 %, v/v) to visualize the brown spots and leaves were photographed by a digital camera. For $O_2^{\cdot-}$ localization *in situ*, samples from each treatment were immersed in a 0.1 mg cm^{-3} solution of nitroblue tetrazolium (NBT) in 50 mM K-phosphate buffer (pH 6.4), containing 10 mM Na-azide, and then were vacuum-infiltrated for 5 - 10 min and left under white light until the appearance of dark spots, characteristic of blue formazan precipitates. The subsequent steps were the same as for localization of H_2O_2 .

One set of cut leaves was placed under HI and periodically taken for Chl fluorescence measurements carried out with an Imaging-PAM Chl fluorometer (Heinz Walz, Effeltrich, Germany) as described by Siebke and Weis (1995) and Rascher *et al.* (2001). All fluorescence measurements were started after 10-min dark adaptation. When performing a measurement, an area of interest (AOI) with a diameter of 1 cm was selected in the middle of the whole leaf. Values of the Chl fluorescence parameters such as variable to maximum fluorescence ratio (F_v/F_m characterising maximum photosystem 2 efficiency), effective photosystem 2 quantum yield ($\Phi_{PS2} = 1 - F_s/F_m'$) and coefficient of photochemical quenching ($qP = (F_m' - F_s)/F_v'$) were determined. In addition, their images were simultaneously derived from the Imaging-PAM software. The definition and calculation of the apparent electron transport rate (ETR) were performed according to Shao *et al.* (2008). Cut leaves placed under HI at 20 °C for 0, 60 and 90 min were also used for rapid light curve measurements (Schreiber *et al.* 1997) carried out using 20-s exposures to stepwise increased irradiance (0, 1, 21, 56, 111, 186, 281, 336, 396, 461, 531, 611, 701, 801, 926, 107, 1251 $\mu\text{mol m}^{-2} \text{s}^{-1}$). Simultaneously, ETR values were obtained automatically using the ImagingWin software.

Reported values of all measurements correspond to the

mean of three to five measurements made on three to five different plants. Student's *t*-test was used to test for statistical differences between two *Arabidopsis* phenotypes. Statistical analyses were performed with

SPSS 12.0 (SPSS, Chicago, IL, USA). Graphing was performed using SigmaPlot 11.0 software (Systat Software, San Jose, CA, USA).

Results

Anthocyanins display a typical absorption peak at 530 nm. The WT showed a marked peak at 530 nm, while the *tt3tt4* anthocyanin-deficient mutant exhibited no obvious peak of absorption at 530 nm. Correspondingly, the anthocyanin content in the WT was nearly four-fold higher than in the *tt3tt4* mutant (data not shown).

After 8 h exposure to HI, there was an obvious decrease in the content of total Chl in the two *Arabidopsis* phenotypes, the *tt3tt4* mutant showing a more pronounced decrease ($P < 0.01$, Fig. 1); however, HI stress had no substantial effect on the carotenoid content. The contents of flavonoids and total phenolics in the two *Arabidopsis* phenotypes were significantly different ($P < 0.001$) prior to HI treatment (Table 1). After 3 - 9 h under HI, the content of flavonoid in the *tt3tt4* mutant decreased by 3 % only, and did not even decrease in the WT. Similarly, the content of total phenolics did not decrease in the both phenotypes. Before HI treatment, no obvious difference was found in the DPPH scavenging capacities between the leaves of *tt3tt4* mutant and WT (Table 1). After HI treatment for 9 h, the DPPH scavenging capacity decreased by 10 % in the *tt3tt4* mutant. By contrast, the DPPH scavenging capacity in the WT increased by 23 %, the increase being mostly related to the increase in the contents of precursors for the synthesis of anthocyanins. Under HI lasting 3 - 9 h, the cell-membrane permeability increased more noticeably in the leaves of the *tt3tt4* mutant as compared to the WT, thus indicating that the plasma membranes in the former suffered more damage (Table 1).

When plants are continuously exposed to HI, ROS accumulate in the cells. Here, H_2O_2 and $O_2^{\cdot-}$ were detected with DAB and NBT staining, respectively, in the leaves of both WT and *tt3tt4* mutant (Fig. 2). In H_2O_2 detection, the brown deposits are the results of the reaction of DAB with

H_2O_2 . In control leaves, some pale brown deposits were observed, without significant differences between WT and *tt3tt4*. As compared to the WT leaf, more brown deposits could be observed in leaves of the *tt3tt4* mutant when both were exposed to HI for 120 min and especially 240 min. In superoxide radical detection, where the blue formazan deposits were characteristic of reaction of NBT with $O_2^{\cdot-}$, the results showed a very similar pattern with the H_2O_2 detection.

In order to identify the levels of photoinhibition, false




Fig. 1. Changes in the contents of chlorophylls (A) and carotenoids (B) in the leaves of two *Arabidopsis* phenotypes (WT: wild type; *tt3tt4*: *tt3tt4* mutant deficient in anthocyanin biosynthesis) exposed to high irradiance ($1\ 300\ \mu\text{mol m}^{-2}\text{s}^{-1}$) for 0 - 8 h. Means \pm SE, $n = 5$.

Table 1. Changes in the contents of total phenolics [$\Delta A_{280}\ \text{mg}^{-1}(\text{f.m.})$] and flavonoids [$\Delta A_{325}\ \text{mg}^{-1}(\text{f.m.})$], DPPH scavenging capacity [$\text{mg}(\text{DPPH})\ \text{g}^{-1}(\text{f.m.})$] and cell-membrane leakage [%] in the leaves of two *Arabidopsis* phenotypes exposed to HI ($1\ 300\ \mu\text{mol m}^{-2}\text{s}^{-1}$) for 0, 3, 6, 9 h. Means \pm SE, $n = 5$. Different letters indicate significant differences between phenotypes at the level $P \leq 0.05$.

Variables	Phenotypes	0 h	3 h	6 h	9 h
Total phenolics	WT	0.10 \pm 0.00a	0.11 \pm 0.00a	0.12 \pm 0.00a	0.11 \pm 0.00a
	<i>tt3tt4</i>	0.09 \pm 0.00b	0.10 \pm 0.00b	0.09 \pm 0.00b	0.10 \pm 0.00b
Flavonoid	WT	0.12 \pm 0.00a	0.13 \pm 0.00a	0.13 \pm 0.00a	0.13 \pm 0.00a
	<i>tt3tt4</i>	0.11 \pm 0.00b	0.12 \pm 0.00b	0.12 \pm 0.00b	0.11 \pm 0.00b
DPPH scavenging capacity	WT	0.31 \pm 0.01a	0.33 \pm 0.01a	0.37 \pm 0.02a	0.38 \pm 0.01a
	<i>tt3tt4</i>	0.29 \pm 0.02a	0.32 \pm 0.02a	0.30 \pm 0.03b	0.26 \pm 0.03b
Membrane leakage	WT	9.25 \pm 0.25a	12.69 \pm 0.21a	15.42 \pm 1.35a	20.50 \pm 1.34a
	<i>tt3tt4</i>	10.38 \pm 0.53a	17.75 \pm 1.23b	29.09 \pm 1.60b	35.23 \pm 2.82b




Fig. 2. H_2O_2 (A) and superoxide radical (B) localization *in situ* in the leaves of two *Arabidopsis* phenotypes (WT and *tt3tt4* mutant). The leaves were exposed to high irradiance ($1\ 300\ \mu\text{mol m}^{-2}\text{ s}^{-1}$) for time indicated, followed by infiltration with DAB or NBT for visualizing H_2O_2 and superoxide radical, respectively.




Fig. 3. Changes in the fluorescence images of the maximal PS 2 quantum yield (F_v/F_m) (A) and effective PS 2 quantum yield (Φ_{PS2}) (B) in the leaves of two *Arabidopsis* phenotypes (WT and *tt3tt4* mutant). Whole leaves were exposed to high-irradiance ($1\ 300\ \mu\text{mol m}^{-2}\text{ s}^{-1}$) for 0 - 180 min. Fluorescence images are indicated by the false colour code at the bottom. The code ranges from black via red, orange, yellow, green, blue and violet to purple, and these colours code for numbers between 0 and 1.




Fig. 4. Effects of high irradiance on the maximum PS 2 quantum yield (F_v/F_m) (A), the electron transport rate (ETR) (B), effective PS 2 quantum yield (Φ_{PS2}) (C) and the coefficient of photochemical quenching (qP) (D) in leaves of two *Arabidopsis* phenotypes (WT: and *tt3tt4* mutant). The whole leaves were exposed to high irradiance ($1300 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 0–180 min. Means \pm SE ($n = 5$).

colour images of F_v/F_m were created in our experiments (Fig. 3A). The distribution of F_v/F_m was not heterogeneous over the unstressed *Arabidopsis* leaf. The mean F_v/F_m value for the whole *Arabidopsis* leaf was about 0.76. After HI treatment for 180 min, the imaging colour of F_v/F_m in the leaves of WT became largely green ($F_v/F_m = 0.4$), although there was a small area of yellow ($F_v/F_m = 0.2$); by contrast, the imaging colour of F_v/F_m in the leaves of *tt3tt4* became yellow ($F_v/F_m = 0.2$), although there was a small area of green ($F_v/F_m = 0.4$). The mean Φ_{PS2} value for the whole *Arabidopsis* leaf was about 0.4 before HI treatment. After HI for 180 min, the imaging colour of Φ_{PS2} in the leaves of WT became partially yellow ($\Phi_{PS2} = 0.2$), although there was a small area of orange ($\Phi_{PS2} = 0.1$). However, the imaging colour in the leaves of *tt3tt4* became partially orange ($\Phi_{PS2} = 0.1$), while there was a small area of black ($\Phi_{PS2} = 0$; Fig. 3B).

Mean values of the Chl fluorescence parameters (F_v/F_m and Φ_{PS2}) from the false colour images were analysed and an obvious decreasing trend in F_v/F_m in *Arabidopsis* leaves was observed during HI treatment (Fig. 4A). After HI for 180 min, the values of F_v/F_m in the leaves of WT and *tt3tt4* decreased by 52.9 and 63.0 % as compared to pre-treatment levels, respectively, which indicated that *tt3tt4* mutant was more sensitive to HI stress than the WT phenotype. In addition, the same pattern also occurred in ETR (Fig. 4B), Φ_{PS2} (Fig. 4C) and qP (Fig. 4D).

The pattern of rapid light curve in the *tt3tt4* mutant was consistent with that of the WT phenotype, and ETR reached a maximum value when the irradiance reached $\sim 1000 \mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 5A). HI stress aggravated the decline of ETR in the two *Arabidopsis* phenotypes and markedly reduced the saturating irradiance of ETR. After treatment for 60 min, the ETR in the leaves of WT and *tt3tt4* reached a saturation value when the irradiance was ~ 1000 and $\sim 900 \mu\text{mol m}^{-2} \text{s}^{-1}$, respectively (Fig. 5B). After

90 min, the saturation irradiance of ETR in the leaves of *tt3tt4* decreased to $\sim 600 \mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 5C).




Fig. 5. Effects of different irradiances and treatment periods on the electron transport rate (ETR) in the leaves of two *Arabidopsis* phenotypes (WT and *tt3tt4* mutant). The treatment was conducted with whole leaves exposed to high irradiance of $1300 \mu\text{mol m}^{-2} \text{s}^{-1}$ for 0, 60 and 90 min, respectively. Then they were exposed (20 s) to stepwise increased irradiance (0, 1, 21, 56, 111, 186, 281, 336, 461, 531, 611, 701, 801, 926, 1076 and $1251 \mu\text{mol m}^{-2} \text{s}^{-1}$). Means \pm SE ($n = 5$).

Discussion

Dihydroflavonol 4-reductase (DFR) is a key enzyme in anthocyanin biosynthesis (Lu *et al.* 2010). The *tt3tt4* mutant of *Arabidopsis*, which is deficient in the DFR locus, exhibited different responses to HI exposure as compared to the WT. It is known that HI often inhibits chlorophyll biosynthesis and increases chlorophyll degradation (Hidema *et al.* 1992, Okada *et al.* 1992, Aarti *et al.* 2007, Zuluaga *et al.* 2008). The total Chl content in the *tt3tt4* mutant showed a more rapid decrease as compared to WT during HI exposure (Fig. 1A, B). This decrease of total Chl was mostly related to the decrease of Chl *a* (data not shown). Malondialdehyde (MDA) is an indicator of lipid peroxidation (Ding *et al.* 2010), which increases membrane leakage. The *tt3tt4* mutant and WT both exhibited increased membrane leakage; however, the former demonstrated a more obvious increase (Table 1). Long-term exposure of a green plant to HI can lead to photoinhibition (Aro *et al.* 1993) and also to production of ROS (Luo *et al.* 2010). It is obvious that the HI-treated leaves of the *tt3tt4* mutant and WT both demonstrated a reduction in PS 2 photochemical efficiency, the former suffering a greater reduction (Figs. 3, 4). In the present study, the *Arabidopsis* leaves were dark-adapted for only 30 min before Chl fluorescence measurement, and thus the mean F_v/F_m of unstressed leaves was only 0.76. It was possible that the unstressed *Arabidopsis* leaves could have a mean F_v/F_m value near 0.80 if they had been dark-adapted overnight. HI stress also led to production of ROS in the two *Arabidopsis* phenotypes, the *tt3tt4* mutant suffering a more severe oxidative stress (Fig. 2). These results indicated that the anthocyanin-deficient mutant of *Arabidopsis* was more susceptible to HI stress as compared to the WT.

We suggest that the higher resistance to HI in the WT was mainly related to the photoprotection provided by anthocyanins. As was mentioned in introduction, the anthocyanins might relieve photooxidative stress by shielding leaf tissues or by quenching ROS. The results of many studies supported the sunscreen hypothesis (*e.g.* Pietrini and Massacci 1998, Close *et al.* 2001, Albert *et al.* 2009). However, some evidence indicates that the optical properties of leaf anthocyanins may not be ideal for a sunscreen role (see review Archetti *et al.* 2009). Moreover, Gould *et al.* (2002b) reported that the effects of the anthocyanins on the penetration of blue and red radiation were negligible. Anthocyanin content in leaves of the WT was low [$0.414 \pm 0.061 \text{ g}^{-1}(\text{f.m.})$; data not shown] and the leaf colour of the WT, the same as the *tt3tt4* mutant deficient in anthocyanin, was green without appearing red. Moreover, there was no significant difference in absorption spectra between the leaves of the two *Arabidopsis* phenotypes (WT and *tt3tt4*) according to our previous research work (Shao *et al.* 2008). Hence, anthocyanins in the WT of *Arabidopsis* could not contribute substantially to shielding during HI stress in the present study. Therefore, we assumed that anthocyanins play an important role possibly as antioxidants rather than

as a sunscreen.

The powerful potential antioxidative capability of anthocyanins has been extensively reported and confirmed by *in vitro* assays (Tsuda *et al.* 1996, Neil *et al.* 2002, Garcia-Alonso *et al.* 2005). A critical point to be considered in conjunction with an antioxidative hypothesis for anthocyanins concerns their localization. Anthocyanin pigments are in the vacuoles of the epidermal cells (Lee and Collins 2001), and there is no report for the presence of anthocyanins in chloroplasts. Although superoxide radicals cannot cross the tonoplast (Takahashi and Asada 1983), they are rapidly protonated to form OH^\bullet or are converted by superoxide dismutase (SOD) to H_2O_2 , which can freely penetrate the tonoplast (Takahashi and Asada 1983, Yamasaki *et al.* 1997). Less accumulation of H_2O_2 and $\text{O}_2^{\bullet-}$ was observed in leaves of the WT compared to the *tt3tt4* mutant (Fig. 2), indicating that anthocyanins can reduce ROS accumulation. In addition, the leaves of two *Arabidopsis* phenotypes showed a similar scavenging capability to DPPH before HI treatment (Table 1). The total antioxidative potential in the anthocyanin-deficient mutant might be compensated by the other antioxidants. During 0–9 h of HI treatment, the DPPH scavenging capacity in WT showed an increasing trend while that in *tt3tt4* showed an increasing trend followed by a decrease. We inferred that the elevated content of anthocyanins led to an increase in DPPH scavenging capacity in the WT. Our previous study reported that anthocyanins *in vitro* were more potent than ascorbic acid (AsA) against DPPH (Zeng *et al.* 2010). Moreover, stress conditions, such as HI, cold temperature, nutrient deficiency or pathogen attack, could lead to accumulations of anthocyanins in plants (*e.g.* Christie *et al.* 1994, Dixon and Paiva 1995, Albert *et al.* 2009). Hence, we suggest that it is possible that the increased precursors for the synthesis of anthocyanins in the WT can result in an increase in DPPH scavenging capacity. Anthocyanins, along with other antioxidants, may protect the leaves of *Arabidopsis* against HI-caused damage, which is consistent with HI tolerance of anthocyanin-rich leaves of a purple rice cultivar (Peng *et al.* 2006).

On the other hand, phenolics and flavonoids, like other antioxidant constituents in plant cells, are also effective ROS scavengers and play an important role in protection against membrane lipid peroxidation in plants (Chalker-Scott 1999, Zhao 1999, Shao *et al.* 2008). They reside in the central vacuole (Hutzler *et al.* 1998), yet their presence in chloroplasts has also been documented (Saunders and McClure 1976). The mutants deficient in enzymes of the anthocyanin biosynthetic pathway, chalcone synthase and dihydroflavonol 4-reductase, may have lower content of phenolics (Shao *et al.* 2008). However, the experimental results of the present study did not confirm it (Table 1).

Previous work in our laboratory confirmed that anthocyanins served as antioxidants to scavenge $\text{O}_2^{\bullet-}$ produced in PS 1 induced by methyl viologen under low

irradiance (Shao *et al.* 2008). In this study, ROS were produced in PS 2 by HI. The results provided strong indirect evidence for the role of anthocyanins in protection against free radicals and their damage to PS 2 in plants. In

conclusion, anthocyanins can be used as antioxidants to extinguish ROS produced in either PS 1 or PS 2 under environmental stresses.

References

- Aarti, D., Tanaka, R., Ito, H., Tanaka, A.: High light inhibits chlorophyll biosynthesis at the level of 5-aminolevulinate synthesis during de-etiolation in cucumber (*Cucumis sativus*) cotyledons. - Photochem. Photobiol. **83**: 171-176, 2007.
- Agati, G., Meyer S., Matteini P., Cerovic Z.G.: Assessment of anthocyanins in grape (*Vitis vinifera* L.) berries using a noninvasive chlorophyll fluorescence method. - J. Agr. Food Chem. **55**: 1053-1061, 2007.
- Albert, N.W., Lewis, D.H., Irving, L.J., Jameson, P.E., Davies, K.M.: Light-induced vegetative anthocyanin pigmentation in *Petunia*. - J. exp. Bot. **60**: 2191-2202, 2009.
- Archetti, M., Döring, T.F., Hagen, S.B., Hughes, N.M., Leather, S.R., Lee, D.W., Lev-Yadun, S., Manetas, Y., Ougham, H.J., Schaberg, P.G., Thomas, H.: Unravelling the evolution of autumn colours: an interdisciplinary approach. - Trends Ecol. Evol. **24**: 166-173, 2009.
- Aro, E.M., Virgin, I., Andersson, B.: Photoinhibition of photosystem II: Inactivation, protein damage and turnover. - Biochim. biophys. Acta **1143**: 113-134, 1993.
- Beggs, C.J., Wellmann, E.: Photocontrol of flavonoid biosynthesis. - In: Kendrick, R.E., Kronenberg, G.H.M. (ed.): Photomorphogenesis in Plants. 2nd Ed. Pp. 733-750. Kluwer Academic Press, Dordrecht – Boston – London 1994.
- Chalker-Scott, L.: Environmental significance of anthocyanins in plant stress responses. - Photochem. Photobiol. **70**: 1-9, 1999.
- Christie, P.J., Alfenito, M.R., Walbot, V.: Impact of low-temperature stress on general phenylpropanoid and anthocyanin pathways: enhancement of transcript abundance and anthocyanin pigmentation in maize. - Planta **194**: 541-549, 1994.
- Close, D.C., Davies, N.W., Beadle, C.L.: Temporal variation of tannins (galloylglucoses), flavonols and anthocyanins in leaves of *Eucalyptus nitens* seedlings: implications for light attenuation and antioxidant activities. - Aust. J. Plant Physiol. **28**: 1-10, 2001.
- De Jong, H.: Inheritance of anthocyanin pigmentation in the cultivated potato: a critical review. - Amer. J. Potato Res. **68**: 585-593, 1991.
- Ding, W., Song, L., Wang, X., Bi, Y.: Effect of abscisic acid on heat stress tolerance in the calli from two ecotypes of *Phragmites communis*. - Biol. Plant. **54**: 607-613, 2010.
- Dixon, R.A., Paiva, N.L.: Stress-induced phenylpropanoid metabolism. - Plant Cell **7**: 1085-1097, 1995.
- Field, T., Lee, D., Holbrook, N.: Why leaves turn red in autumn. The role of anthocyanins in senescing leaves of red-osier dogwood. - Plant Physiol. **127**: 556-574, 2001.
- Fukumoto, L.R., Mazza, G.: Assessing antioxidant and prooxidant activities of phenolic compounds. - J. Agr. Food Chem. **48**: 3597-3604, 2000.
- Garcia-Alonso, M., Rimbach, G., Sasai, M., Nakahara, M., Matsugo, S., Uchida, Y., Rivas-Gonzalo, J.C., De Pascual-Teresa, S.: Electron spin resonance spectroscopy studies on the free radical scavenging activity of wine anthocyanins and pyranoanthocyanins. - Mol. Nutr. Food Res. **49**: 1112-1119, 2005.
- Gould, K.S., Kuhn, D.N., Lee, D.W., Oberbauer, S.F.: Why leaves are sometimes red. - Nature **378**: 241-242, 1995.
- Gould, K.S., Lister, C.: Flavonoid functions in plants. - In: Andersen, O.M., Markham, K.R. (ed.): Flavonoids: Chemistry, Biochemistry and Applications. Pp. 397-411. CRC Press, Boca Raton 2006.
- Gould, K.S., McKelvie, J., Markham, K.R.: Do anthocyanins function as antioxidants in leaves: imaging of H₂O₂ in red and green leaves after mechanical injury. - Plant Cell Environ. **25**: 1261-1269, 2002a.
- Gould, K.S., Vogelmann, T.C., Han, T., Clearwater, M.J.: Profiles of photosynthesis within red and green leaves of *Quintinia serrata* A. - Cunn. Physiol. Plant. **116**: 127-133, 2002b.
- Havaux, M., Niyogi, K.K.: The violaxanthin cycle protects plants from photooxidative damage by more than one mechanism. - Proc. nat. Acad. Sci. USA **96**: 8762-8767, 1999.
- Hidema, J., Makino, A., Kurita, Y., Mae, T., Ojima, K.: Changes in the levels of chlorophyll and light-harvesting chlorophyll a/b protein of PSII in rice leaves aged under different irradiances from full expansion through senescence. - Plant Cell Physiol. **33**: 1209-1214, 1992.
- Hoch, W.A., Zeldin, E.L., McCown, B.H.: Physiological significance of anthocyanins during autumnal leaf senescence. - Tree Physiol. **21**: 1-8, 2001.
- Huner, N.P.A., Öquist, G., Sarhan, F.: Energy balance and acclimation to light and cold. - Trends Plant Sci. **3**: 224-230, 1998.
- Hutzler, P., Fischbach, R., Heller, W., Jungblut, T.P., Reuber, S., Schmitz, R., Veit, M., Weissenböck, G., Schnitzler, J.P.: Tissue location of phenolic compounds in plants by confocal laser scanning microscopy. - J. exp. Bot. **49**: 953-965, 1998.
- Ismail, G.S.M., Mohamed, H.E.: Alteration in growth and thylakoid membrane lipid composition of *Azolla caroliniana* under phosphate deficiency. - Biol. Plant. **54**: 671-676, 2010.
- Lee, D.W., Collins, T.M.: Phylogenetic and ontogenetic influences on the distribution of anthocyanins and betacyanins in leaves of tropical plants. - Int. J. Plant Sci. **162**: 1141-1153, 2001.
- Lichtenthaler, H.K., Wellburn, A.R.: Determinations of total carotenoids and chlorophyll *a* and *b* in leaf extracts of different solvents. - Biochem. Soc. Trans. **603**: 591-592, 1983.
- Lu, X., Zhou, W., Gao, F.: Chromosomal location of 45S rDNA and dfr gene in *Citrus sinensis*. - Biol. Plant. **54**: 798-800, 2010.
- Luo, Y., Li, F., Wang, G.P., Yang, X.H., Wang, W.: Exogenously-supplied trehalose protects thylakoid membranes of winter wheat from heat-induced damage. - Biol. Plant. **54**: 495-501, 2010.
- Manetas, Y.: Why some leaves are anthocyanic and why most anthocyanic leaves are red. - Flora **201**: 163-177, 2006.
- Manetas, Y., Petropoulou, Y., Psaras, G.K., Drinia, A.: Exposed red (anthocyanic) leaves of *Quercus coccifera* display shade characteristics. - Funct. Plant Biol. **30**: 265-270, 2003.

- Meng, X.C., Xing, T., Wang, X.J.: The role of light in the regulation of anthocyanin accumulation in *Gerbera hybrida*. - Plant Growth Regul. **44**: 243-250, 2004.
- Merzlyak, M.N., Chivkunova, O.B., Solovchenko, A.E., Naqvi, K.R.: Light absorption by anthocyanin in juvenile, stressed, and senescing leaves. - J. exp. Bot. **59**: 3903-3911, 2008.
- Neil, S., Gould, K.S., Kilmartin, P.A., Mitchell, K.A., Markham, K.R.: Antioxidant activities of red versus green leaves in *Elatostema rugosum*. - Plant Cell Environ. **25**: 539-547, 2002.
- Neill, S.O., Gould, K.S.: Anthocyanins in leaves: light attenuators or antioxidant. - Funct Plant Biol. **30**: 865-873, 2003.
- Okada, K., Inoue, Y., Satoh, K., Katoh, S.: Effects of light on degradation of chlorophyll and proteins during senescence of detached rice leaves. - Plant Cell Physiol. **33**: 1183-1191, 1992.
- Peer, W.A., Brown, D.E., Tague, B.W.: Flavonoid accumulation patterns of transparent testa mutants of *Arabidopsis*. - Plant Physiol. **126**: 536-548, 2001.
- Peng, C.L., Chen, S.W., Lin, Z.F., Lin, G.Z.: [Detection of antioxidative capacity in plants by scavenging organic free radical DPPH.] - Progr. Biochem. Biophys. **27**: 658-661, 2000. [In Chin.]
- Peng, C.L., Lin, Z.F., Lin, G.Z., Chen, S.W.: The anti-photooxidation of anthocyanin-rice leaves of a purple rice cultivar. - Sci. China Ser. C **49**: 543-551, 2006.
- Pietrini, F., Massaei, A.: Leaf anthocyanin content changes in *Zea mays* L. grown at low temperature: significance for the relationship between quantum yield of PS II and the apparent quantum yield of CO₂ assimilation. - Photosynth. Res. **58**: 213-219, 1998.
- Piovan, A., Filippini, R.: Anthocyanins in *Catharanthus roseus* *in vivo* and *in vitro*: a review. - Phytochem. Rev. **6**: 235-242, 2007.
- Rascher, U., Hütt, M.T., Siebke, K., Osmond, B., Beck, F., Lüttge, U.: Spatio-temporal variation of metabolism in a plant circadian rhythm: the biological clock as an assembly of coupled individual oscillators. - Proc. nat. Acad. Sci. USA **98**: 11801-11805, 2001.
- Romero-Puertas, M.C., Rodriguez-Serrano, M., Corpas, F.J., Gómez, M., Delrío, L.A., Sandalio, L.M.: Cadmium-induced subcellular accumulation of O₂⁻ and H₂O₂ in pea leaves. - Plant Cell Environ. **27**: 1122-1134, 2004.
- Saunders, J.A., McClure, J.W.: Distribution of flavonoids in chloroplasts of 25 species of vascular plants. - Phytochemistry **15**: 809-810, 1976.
- Schreiber, U., Gademann, R., Ralph, P.J., Larkum, A.W.D.: Assessment of photosynthetic performance of prochloron in *Lissoclinum patella* by in situ and in hospite chlorophyll fluorescence measurements. - Plant Cell Physiol. **38**: 945-951, 1997.
- Shao, L., Shu, Z., Sun, S.L., Peng, C.L., Lin, Z.F., Yang, C.W.: Enhanced sensitivity of *Arabidopsis* anthocyanin mutants to photooxidation: a study with fluorescence imaging. - Funct. Plant Biol. **35**: 714-724, 2008.
- Shirley, B.W., Qubasek, W.L., Storz, G., Bruggemann, E., Koornneef, M., Ausubel, F.M., Goodman, H.M.: Analysis of *Arabidopsis* mutants deficient in flavonoid biosynthesis. - Plant J. **8**: 659-671, 1995.
- Siebke, K., Weis, E.: Assimilation images of leaves of *Glechoma hederacea*: analysis of nonsynchronous stomata related oscillations. - Planta **196**: 155-165, 1995.
- Smillie, R.M., Hetherington, S.E.: Photoabatement by anthocyanin shields photosynthetic systems from light stress. - Photosynthetica **36**: 451-463, 1999.
- Steyn, W.J., Wand, S.J.E., Holcroft, D.M., Jacobs, G.: Anthocyanins in vegetative tissues: a proposed unified function in photoprotection. - New Phytol. **155**: 349-361, 2002.
- Takahashi, M.A., Asada, K.: Superoxide anion permeability of phospholipid- membranes and chloroplast thylacoids. - Arch. Biochem. Biophys. **226**: 558-566, 1983.
- Tsuda, T., Shiga, K., Ohshima, K.: Inhibition of lipid peroxidation and the active oxygen radical scavenging effect of anthocyanin pigment isolated from *Phaseolus vulgaris* L. - Biochem. Pharmacol. **52**: 1033-1039, 1996.
- Wade, H.K., Sohal, A.K., Jenkins, G.I.: *Arabidopsis* ICX1 is a negative regulator of several pathways regulating flavonoid biosynthesis genes. - Plant Physiol. **131**: 707-715, 2003.
- Wang, H., Cao, G.H., Prior, R.L.: Oxygen radical absorbing capacity of anthocyanins. - J. Agr. Food Chem. **45**: 304-309, 1997.
- Yamasaki, H., Sakihama, Y., Ikehara, N.: Flavonoid-peroxidase reaction as a detoxification mechanism of plant cells against H₂O₂. - Plant Physiol. **115**: 1405-1512, 1997.
- Zeng, X.Q., Chow, W.S., Su, L.J., Peng, X.X., Peng, C.L.: Protective effect of supplemental anthocyanins on *Arabidopsis* leaves under high light. - Physiol. Plant. **138**: 215-225, 2010.
- Zhao, B.L.: [Oxygen free radicals and natural antioxidants]. - Science Press, Beijing 1999. [In Chin.]
- Zuluaga, D.L., Gonzali, S., Loreti, E., Pucciariello, C., Degl'Innocenti, E., Guidi, L., Alpi, A., Perata, P.: *Arabidopsis thaliana* MYB75/PAP1 transcription factor induces anthocyanin production in transgenic tomato plants. - Funct. Plant Biol. **35**: 606-618, 2008.