

Table 1 Suppl. Forward (F) and reverse (R) primers used for reverse transcription quantitative PCR assays.

Gene	Primer sequence
Modified <i>MxProtox</i>	F: ACAGGATGAGGATGCTTTGG R: CTGCTAGTTGAGCTGCGTTG
<i>OsProtox</i>	F: ACAGTTCCTCATTGGCCATC R: CCCATGAAATTTTGTGCT
<i>FC2</i>	F: TTGGTGCTATGGCAGTTTCA R: AGTGGAACAAAGGCAGGATG
<i>HO2</i>	F: AGGGACCTAGCAGCCCTAAC R: CCCGTATCGTCCATCTTGAG
<i>CatB</i>	F: AAGGATGGAAAGCCTCACCT R: AGGGGGATGATATCCTCTGG
<i>CatC</i>	F: ACAACCACTACGACGGCTTC R: CCAGTAGGAGAGCCAGATGC
<i>APXa</i>	F: ACAAAGCCCTGCTGAGTGAC R: TAACAGCCCACCGAGACATT
<i>APXb</i>	F: CCAAGTGACAAAGCCCTCAT R: TCTTGACAGCAAATAGCTTGG
<i>Actin</i>	F: CTTCATAGGAATGGAAGCTGCGGGTA R: CGACCACCTTGATCTTCATGCTGCTA