

Differential distribution of transcripts from genes involved in polyamine biosynthesis in bean plants

J.F. JIMÉNEZ-BREMONT¹, E. HERNÁNDEZ-LUCERO, A.G. ALPUCHE-SOLÍS, S. CASAS-FLORES and A.P. BARBA DE LA ROSA

División de Biología Molecular, Instituto Potosino de Investigación Científica y Tecnológica, Camino a la Presa de San José 2055, 78210 Tangamanga, San Luis Potosí, SLP, México

Abstract

Partial cDNAs sequences for arginine decarboxylase (*Pvadc*), S-adenosylmethionine decarboxylase (*Pvsamdc*) and spermidine synthase (*Pvspds*) were isolated from the bean *Phaseolus vulgaris* using primers designed from conserved regions of enzymes belonging to plant species. Sequence analysis showed that the *Pvadc*, *Pvsamdc* and *Pvspds* genes were most closely related to the orthologous genes from *Glycine max*, *Phaseolus lunatus* and *Pisum sativum*, respectively. The expression patterns of the genes, together with that of ornithine decarboxylase (*Pvodc*), were analysed in young and mature leaves, stems, roots, root tips, petals, stigma, ovaries, filaments and anthers of bean plants. *Pvsamdc* was found to be expressed at similar levels in all tissues. The other transcripts showed tissue specific expression. *Pvadc* was barely expressed in petals and not at all in roots tips, *Pvspds* was mainly expressed in roots, stigma and filaments, and *Pvodc* was detected only in roots.

Additional key words: arginine decarboxylase, ornithine decarboxylase, *Phaseolus vulgaris*, S-adenosylmethionine decarboxylase, spermidine synthase.

Introduction

Polyamines are essential and abundant organic cations known to be involved in many basic processes in plants (Tiburcio *et al.* 1990, Kumar *et al.* 1997, Malmberg *et al.* 1998). The most abundant polyamines are putrescine (Put), spermidine (Spd) and spermine (Spm). Polyamines (free or conjugated) have been related specifically to the response to stress, rhizogenesis (Jarvis *et al.* 1985, Kakkar *et al.* 2000), root development and meristematic function, somatic embryogenesis and organogenesis (Feirer *et al.* 1984), flower and fruit development (Kakkar and Rai 1993, Galston and Kaur-Sawheny 1990), pollen formation and dormance (Kakkar and Nagar 1997), growth response induced by light (Yoshida *et al.* 1999), lateral growth of leaf-homolog organs (Watson *et al.* 1998), senescence (Muchitch *et al.* 1983, Kakkar and Nagar 1996), and response to microbial symbionts (El Ghachoui *et al.* 1996). The first polyamine synthesized in the metabolic route is Put, derived from ornithine through the activity of ornithine decarboxylase (ODC; EC 4.1.1.17). In plants and some bacteria, Put is

also synthesized from arginine by the action of arginine decarboxylase (ADC; EC 4.1.1.19). Put is subsequently converted to Spd through the activity of S-adenosylmethionine decarboxylase (SAMDC; EC 4.1.1.50) and spermidine synthase (SPDS; EC 2.5.1.16). Conversion of Spd to Spm also requires SAMDC and spermine synthase (SPMS; EC 2.5.1.22) (Tabor and Tabor 1984).

Expression patterns of the gene encoding ADC support the idea that this is the main source of putrescine during cell elongation. Studies on ODC suggest that it may be involved in regulation of cell division in growing tissues (Michael *et al.* 1996, Nam *et al.* 1997, Martin-Tanguy 1997, Alabadi and Carbonell 1998). Several studies have shown that the *samdc* transcript is mainly accumulated in young tissues and organs where active cell division occurs, and decline to lower levels as organs age and senesce (Mad Arif *et al.* 1994, Marco and Carrasco 2002). In contrast, the *spds* transcript is expressed in all plant organs in *Arabidopsis thaliana* (Hanzawa *et al.* 2002). Plant *samdc*s possess a small upstream ORF

Received 23 September 2004, accepted 1 September 2005.

Abbreviations: ADC - arginine decarboxylase; ODC - ornithine decarboxylase; ORF - open reading frame; PCR - polymerase chain reaction; Put - putrescine; RT-PCR - reverse-transcriptase polymerase chain reaction; SAMDC - S-adenosylmethionine decarboxylase; Spd - spermidine; SPDS - spermidine synthase; Spm - spermine; SPMS - spermine synthase.

Acknowledgements: We thank Dr. José Ruiz-Herrera and Dr. Rogerio Sotelo Mundo for critical review of the manuscript.

¹ Corresponding author; fax (+52) 444 8342010, e-mail: jbremont@ipicyt.edu.mx.

(uORF) in the 5' leader sequence of the mRNA. Hanfrey *et al.* (2002) demonstrated that uORF-mediated translational control of SAMDC is essential for polyamine homeostasis and for normal growth and development.

Previously, we have reported the ODC-encoding gene (*Pvodc*) in *Phaseolus vulgaris* (Jiménez-Bremont *et al.*

2004). The present communication describes the isolation and sequencing of partial cDNA sequences encoding ADC, SPDS, SAMDC and the *samdc* uORF from *Phaseolus vulgaris*. Based on the isolated gene sequences, specific primers were designed in order to analyse gene expression patterns in different bean tissues.

Materials and methods

Plants: *Phaseolus vulgaris* L. (cv. Canario 60) seeds were sterilized with a 50 % commercial sodium hypochlorite solution (6 % free chlorine) for 10 min and rinsed several times with sterile, distilled water. Aseptic seeds were germinated in an agrolite/water support, and grown in a commercial substrate (*Cosmocel*, Monterrey, Mexico) in plastic pots for 45 d. Plants were maintained until flowering in a growth chamber under controlled conditions (16-h photoperiod, irradiance 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$, temperature $25 \pm 2^\circ\text{C}$).

RNA isolation and RT-PCR assays: Total RNA was isolated using the *Trizol* reagent as recommended by the manufacturer (*Invitrogen*, Carlsbad, USA), and treated with DNase I (*Invitrogen*). First strand cDNA synthesis was performed with SuperScriptTM First-Strand Synthesis System for reverse-transcriptase polymerase chain reaction (RT-PCR; *Invitrogen*). Reaction mixtures were incubated with RNase H at 42 °C for 50 min, then at 70 °C for 15 min, and finally at 37 °C for 20 min to remove RNA templates from the cDNA:RNA hybrid molecules. A control reaction for each sample was performed without reverse transcriptase. The samples were analysed by electrophoresis on 1 % agarose gels. PCR products of the expected size were cloned in the pCR 2.1 vector (*TA Cloning Kit, Invitrogen*) and further sequenced. Alignments and phylogenetic analyses were performed with the MegAlign (*DNASTAR*, Madison, USA) and *Clustal* (Higgins and Sharp 1988) programs. Sequences were compared to those available at EMBL GenBank.

Cloning and sequencing of *Pvadc*, *Pvspds*, *Pvsamdc* and uORF_{samdc} cDNAs fragments: Primers (external or internal) of each gene were designed, based on conserved regions of the corresponding enzymes from several plants. Nested PCR reactions were started with the external primers followed by further amplification using the internal primers. The template was the cDNA obtained by reverse transcription using RNA isolated from bean leaves. The primers synthesized for the isolation of the *Pvadc* fragment were: 38 5'-CTCA TTACCAAGGTGTTATCCGGTGAA-3' (sense), 40 5'-GAGCGCCTCCTCATAAGCCCCACCCAAA-3' (antisense) and 42 5'-GGCTTCTCGTCCAAACGG TGAAT-3' (antisense) designed according to the conserved regions of cDNAs from: *Glycine max* (GenBank U35367), *Vitis vinifera* (GenBank X96791),

Nicotiana tabacum (GenBank AF321137.1) and *Datura stramonium* (GenBank AJ251898.1). Primers to obtain the *samdc* fragment were: 2272 5'-GTTTCTGCAATT GGTTTGAAAGGTTT-3' (sense), 2274 5'-GAAAAGA GGTTGGAAATATCCT-3' (sense), 2273 5'-CTTCTT CATCTTCATCTTCCAGCA-3' (antisense) and 2299 5'-CCACCCATTCCAAGC-3' (antisense) from conserved regions of cDNAs from: *Phaseolus lunatus* (GenBank AB062360), *G. max* (GenBank AF488307), *Pisum sativum* (GenBank AB087841.1) and *Arabidopsis thaliana* (GenBank U63633). Primers for synthesis of the *Pvspds* fragment were: Spe-3-5 5'-CCTGGATGGTT CTCTGAAATTAGCC-3' (sense) and Spe-3-3 5'-CTC TTCGCAAAAGATGGCAAACAGAA-3' (antisense) from conserved regions of cDNAs from: *P. sativum* 2 (GenBank AF043109.1), *P. sativum* 1 (GenBank AF043108), *Lycopersicon esculentum* (GenBank AJ006414) and *D. stramonium* 2 (GenBank Y08253). For the uORF isolation, we designed the primer uORF 5'-ATGGAGTCTAAAGGAGGTAAAAAG-3' (sense) located at the beginning of the small conserved open reading frame in the 5' UTR found in plants *samdc*'s (Franceschetti *et al.* 2001) in combination with primer 2299 5'-CCACCCATTCCAAGC-3' (antisense) (Fig. 3A).

RT-PCR analyses of gene transcripts: Different sets of primers were designed to amplify each cDNA by RT-PCR: *Pvodc* (GenBank AY125817) sense 5'-ATGTTGG CTGCTATGGGTTCGAA-3' and antisense 5'-AGTAT TCACGTAAATCTCCACGTA-3'. *Pvadc* (GenBank AY671973) sense RTADC1 5'-TCGTGGTAGAGG ACATCGTAAA-3' and antisense RTADC2 5'- CCAA ACGGTGAATGGGGATGA-3'; *Pvsamdc* (GenBank AY327898) sense RTSAM1 5'-GGTTTAAGA GTTCTTACGAAATCAC-3' and antisense RTSAM2 5'- ATTTGTTCAGAGGGCTTGCTTGC-3'; *Pvspds* (GenBank AY674166) sense RTSPE1 5'-TGATCA CTCATCTTCCTCTTGCTC-3' and antisense RTSPE2 5'-CTCAGAGTTGAAAATTCAGTGGTC-3'. For synthesis of an actin fragment (loading control) the primers used were 5-Act 5'-ATGGGGCAGAAGGAT GCGTATG-3' (sense) and 3-Act 5'-AGCCTTCA TAGATGGGGACCGT-3' (antisense). Equal RNA amounts were used for each RT-PCR reaction as described previously. PCR amplifications were performed in a 0.05 cm³ reaction mixture containing 0.001 cm³ of the RT reaction product as template. The PCR amplification conditions for each gene were

as follows: 5 min at 94 °C, 30 cycles of 45 s at 94 °C, 45 s at 55 °C (*Pvodc*), 56 °C (*Pvadc*), 50 °C (*Pvsamdc*) or

58 °C (*Pvspds*), 80 s at 72 °C and 8 min at 72 °C for the final extension.

Results and discussion

Plant ADC, SAMDC and SPDS sequences have highly conserved regions and thus provide several candidate sites for designing PCR primers of consensus sequence in order to clone related genes. Nested amplifications by PCR were carried out using these primers to obtain partial cDNAs sequences encoding ADC, SAMDC and SPDS from *Phaseolus vulgaris*.

Isolation and sequence analysis of a partial cDNA corresponding to a bean arginine decarboxylase gene (*Pvadc*): We amplified a unique band of 1161 bp that corresponded to the gene encoding *P. vulgaris* ADC. The sequence obtained (Fig. 1A) was registered at the GenBank (AY671973). The pyridoxal phosphate binding site (Sandmeier *et al.* 1994), was identified at amino acids

7 - 25 [YP(V)KCN(QDRF)VV(ED)I(VKF)G] of the encoded sequence (Fig. 1A). Site-directed mutagenesis of Lys95 in *Nicotiana glutinosa* ODC causes a substantial loss of enzymatic activity (Lee and Cho 2001). The consensus sequence of the decarboxylases family responsible for substrate binding was located at amino acids 187 - 201 [G(AN)M(RV)IDIGGGLG] of the *P. vulgaris* encoded sequence. It has been proposed that the stretch of three consecutive glycines is the motif directly involved in substrate-binding (Moore and Boyle 1990). A probable PEST region [(sequences rich in proline (P), glutamic acid (E), serine (S) and threonine (T)] characteristic of proteins with high turnover rate (Rechsteiner and Rogers 1996), with PEST score of -8.45, was identified at residues 255 - 292 of the encoded

A) ADC

```
Pv -----
Gm MPVLACCVDAAAPPGYAFAGDISFPAPIAFTGVPPATADDTNNSNNHWSPSLSAALYINVGDWGPPYFAVNTAGNISVRPH 80
At 2 MPALACVDTSEVPAYAFS-----DTAGDVFIPASSPTSAAVVUDRWSPSLSSSIYRLTDGWGA PYFTANSSGNTISVRPH 74
```

Pv	-----	Primer 38	SHYQGVYPVKCNQDRFV	17
Gm	GSDTVSHQEIDLKIVKKASDPKSLGGQLQLPLIARFPDVLRNRLQSAFDYAIQSGGYE	SHYQGVYPVKCNQDRFV	160	
At2	GSETLPHQDIDLKIVKKVTGPKSSGGLQLPLI	RVFPDVLRNRLQSAFDYAIKSQGYD	SHYQGVYPVKCNQDRFV	154

Pv	KKLCIKPVIGLRAKLRKHS G HGGYFRRESKFG L TTA Q ILRV V KKL D LAGM L DC L QLL H FHIGS Q IP S TA L ADGV E A	177
Gm	KKLCIKPVIGLRAKLRKHS G HGGI F R R K G F L TTAR V LR V KKN L DLAGM L DC L QLL H FHIGS Q IP S TA L ADGV E A	320
At 2	QKMNVRPVIGLRAKLRKHS G HGF G STSGE K GK F LTT Q IV V VR V R L RQSGM L DC L QLL H FHIGS Q IP S TS L SDGV A EEA	314
***	*****	*****

Pv	AQIYCELVRIGANMRVIDIGGGGL	IDYDGSKSCDSDISVGYSLEEYASAVVHAIQCVCDDR SVKNPVICS ESGRAIVSHQ	257
Gm	AQIYCELVRIGANMRVIDIGGGGL	IDYDGSKSCDSDISVEYGLEEYAAAVVHAVQCVC DR -SVKHPVICS ESGRAIVSHH	400
At 2	AQLYCELVRIGAHMKVVIDIGGGGL	IDYDGSKSGESSDLSVASYLEEYAAEVVASVRVVC DRSSVKHPVICS ESGRAIVSHH	394

Pv	SILVFEAVGTSTVGG-VSSAFA S A P Y V A G - - D L S E D Y R F L S E A F G G D Y E S C L A Y T E E L K E R C V E Q F K Q G A V C M E Q L A A V	336
Gm	S V L I F E A V G T S S T N G G G A P P A L S A H Y L A E - - E L S E D Y G Y L S E L A F R G D Y E T C L V Y T E E M K E R C V E Q F K Q G T V C M E Q L A A V	479
At 2	S V L I F E A V S A D K P M V H Q A T P C D I Q F L L E G N E E A R A N Y E D L Y A V M R G D H E S C L L Y V D Q L K Q R C V E G F K E G V L S I E Q L A S V	474

Pv	DGLCELVRKAVGAGEPVRRYHVNLSVFTSIPDAWGLIEQVFPIIPIHLRLEDKP-----	400
Gm	EGLCELVRKAVGAAESVRRYHVNLISFTSPDAWGLIEQVFPIIPIHLRLEDKPSVRGILSDLTCDSGKIDKFINGESSLP	557
At 2	DGLCEWVLKAIGASDPVHTYNIINLNSVFTSIPDLWGLIDQLFPIVPIHKLDQRPGARGILSDLTCDSGKINKFIGGESSLP	557

Pv -----
 Gm LH~~ME~~ME--GGRT~~Y~~YLG~~M~~FLGGAYEEALGGVHNL~~F~~GGPSVV~~R~~VSQSDGPHSFAVTRAVPGPSCGDV~~L~~VRM~~Q~~HQPE~~L~~MF~~T~~L 634
 At 2 LH~~E~~LDKNGSGGRYFLG~~M~~FLGGAYEEALGGVHNL~~F~~GGPSVV~~R~~VSQSDGPHSFAVTRAVPGQSSADVL~~R~~AMQHEPE~~L~~MF~~T~~L 634

B) SPDS

C) SAMDC

Fig. 1. Comparison of predicted amino acid sequences encoded by the cDNA fragments of *Phaseolus vulgaris* ADC, SAMDC and SPDS with the corresponding enzymes from different plants. A) ADC: *P. vulgaris*, *Glycine max* and *Arabidopsis thaliana* 2. The consensus sequence of the decarboxylases family is shown in a box; the piridoxal binding site is underlined; and double underlining marks the possible PEST regions. B) SPDS: *P. vulgaris*, *Pisum sativum* 2 and *A. thaliana*. The consensus sequence of the synthases family appears in a box; the six proposed binding motifs for dcSAM are gray-highlighted; and a probable PEST region is double underlined. C) SAMDC: *P. vulgaris*, *P. lunatus* and *A. thaliana*. The consensus sequence of the decarboxylases family and the site of proenzyme processing are shown in a box; and the probable PEST regions are shown in lowercase. Identical residues are marked by asterisks, and amino acid conserved substitutions are indicated as dots.

sequence (Fig. 1A). Comparison using *Clustal* analysis of this fragment with other plant ADCs showed an homology of 87.6 % with *Glycine max* (GenBank U35367), 73.6 % with *Vitis vinifera* (GenBank X96791), 73.1 % with *Nicotiana tabacum* (GenBank AF321137.1),

72.6 % with *Datura stramonium* (GenBank AJ251898.1) and 72.4 % with *Capsicum annuum* (GenBank AY156514.1). According to these results, bean ADC is most closely related to the *G. max* enzyme (Fig. 2).

Isolation and sequence analysis of a partial cDNA corresponding to a bean spermidine synthase gene (*Pvspds*): We isolated by PCR a single band (728 bp; GenBank AY674166) corresponding to a cDNA fragment of *Pvspds* from bean. The encoded sequence showed high similarity with other SPDSS: 88.5 % with *Pisum sativum* 2 (GenBank AF043109.1), 87.2 % with *P. sativum* 1 (GenBank AF043108), 84.1 % with *Lycopersicon esculentum* (GenBank AJ006414), 82.3 % with *D. stramonium* 2 (GenBank Y08253), 81.4 % with *Arabidopsis thaliana* (GenBank AB025622.1) (Fig. 2).




Fig. 2. Phylogenetic dendograms of ADCs, SPDSs and SAMDCs. ADC: *Datura stramonium*, *Nicotiana tabacum*, *Lycopersicon esculentum*, *Capsicum annuum*, *Nicotiana sylvestris*, *Arabidopsis thaliana* 1, *A. thaliana* 2, *Carica papaya*, *Vitis vinifera*, *Malus domestica*, *Daucus carota*, *Phaseolus vulgaris*, *Glycine max*, *Pisum sativum* and *Avena sativa*. SPDS: *D. stramonium* 2, *Hyoscyamus niger*, *L. esculentum*, *Solanum tuberosum*, *C. annuum*, *N. sylvestris*, *N. tabacum*, *D. stramonium* 1, *Oryza sativa*, *P. sativum* 2, *P. vulgaris*, *P. sativum* 1, *M. domestica*, *G. max*, and *A. thaliana*. SAMDC: *Phaseolus lunatus*, *P. vulgaris*, *G. max*, *P. sativum*, *Vicia faba*, *A. thaliana*, *D. carota*, *N. tabacum*, *V. vinifera*, *M. domestica*, *Magnifera indica*, *Ipomoea nil*, *Oryza sativa*, and *Zea mays*.

A consensus sequence of the synthases family [VLVIGGG(D)G(GV)L(R)E] (Sekowska *et al.* 1998), was located at amino acids 37 - 51 of the encoded sequence (Fig. 1B). Six SAM or dcSAM-binding motifs were identified in the sequence. Motif I [IGGGDGGVLRREIARHSS] corresponding to amino acids 40 - 57, shares 11 amino acids with the consensus sequence of the synthases family, motif II [EI] is located at amino acids 65 - 66, motif III [DG] at amino acids 96 - 97, motif IV [DAVIVDSS] at amino acids 110 - 117, motif V [GPAQELF] at amino acids 121 - 127 and motif VI [PGGVVCTQAA] is located at amino acids 141-149 (O'Gara *et al.* 1995) (Fig. 1B). Also, a possible PEST region with a score of -1 was located at amino acids 102 - 129 [KNVPEGTYDAVIVDSSDPIGPAQELFEK] (Fig. 1B).

Isolation and sequence analysis of a partial cDNA corresponding to a bean S-adenosylmethionine decarboxylase gene (*Pvsamdc*): The nested PCR assays produced a single fragment of 935 bp. We compared the sequence of the polypeptide encoded by *P. vulgaris samdc* (GenBank AY327898, reported in this work) with the sequences reported from other plant SAMDCs: The highest similarity was to *P. lunatus* with 95.2 % (GenBank AB062360) and to *G. max* with 82.3 % (GenBank AF488307) enzymes; lower homology to enzymes from *P. sativum* with 75.6 % (GenBank AB087841.1), *Vicia faba* with 74.6 % (GenBank CAB76966) and *A. thaliana* with 66.6 % (GenBank U63633), was shown; whereas sequence comparison with enzymes from the monocots *Oryza sativa* (GenBank Y07766) and *Zea mays* (GenBank Y07767) SAMDCs showed 53.7 % and 51.4 %, respectively. These results suggest that the homology of SAMDCs agrees with the phylogenetic relationships of the plant species (Fig. 2). The putative characteristic motifs of decarboxylases were identified in the encoded sequence (Fig. 1C). Accordingly, a conserved sequence from all SAMDCs was located at amino acid residues 50 - 61 of the encoded sequence [SYVLSESSLFV], which is also considered as the consensus proenzyme-processing site of the decarboxylases family (Ekstrom *et al.* 1999). Two putative PEST regions located at amino acids 26 - 65 [KSQLDEILTPAACTIVSSLSNDDVDSYVLSESSLFV YAYK] and 234-254 [HITPEDGFSYASFETVGYDFK] of the encoded sequence with scores of -6.52 and -7.17 respectively, were identified (Fig. 1C). Interestingly, the first putative PEST box is located at the proenzyme-processing site of the enzyme (Mad Arif *et al.* 1994).

5' UTR ORF in *Pvsamdc* of *P. vulgaris*: A single 1316 pb fragment, which includes the bean *samdc* 5' UTR region and a small ORF within this sequence was amplified by PCR. The sequence obtained corresponding to the 5' UTR ORF from *P. vulgaris samdc* and was registered at the GenBank (AY730047). As indicated above, the encoded small uORF (53 aa) has been described to be involved in the translational control of




Fig. 3. Alignment of the deduced amino acid sequences of the small uORF harbored in the *samdc* mRNA 5' leader sequences. A) Scheme of the strategy followed to obtain the uORF sequence; see text for details. B) EST sequences of several monocots and dicots were compared (accession numbers in parentheses): Le, *Lycopersicon esculentum* (EST-AI483261); Ds, *Datura stramonium* (Y07768); Pt, *Pinus taeda* (EST-AI725223); Nt, *Nicotiana tabacum* (AF033100); Cl, *Citrullus lanatus* (EST-AI563097); Gm, *Glycine max* (EST-AI442381); Gh, *Gossypium hirsutum* (EST-AI728571); Cr, *Catharanthus roseus* (U12573); In, *Ipomoea nil* (U64927); Pv, *Phaseolus vulgaris* (AY730047); Ps, *Pisum sativum* (U60592); Mt, *Medicago trunculata* (EST-AA661003); At2, *Arabidopsis thaliana* 2 (AJ251899); Bj2, *Brassica juncea* 2 (X95729); Bj1, *Brassica juncea* 1 (U80916); Zm, *Zea mays* (Y07767); Hc, *Hordeum chilense* × *Triticum turgidum* conv. Durum (X83881); Os1, *Oryza sativa* 1 (Y07766); Os2, *Oryza sativa* 2 (AJ251899); Dc1, *Dianthus caryophyllus* (U38526); Dc2, *Dianthus caryophyllus* (U38527); Mc, *Mesembryanthemum crystallinum* (EST-AI822872). Identical residues are marked by asterisks, and amino acid conserved substitutions are indicated as dots.

SAMDC in plants (Hanfrey *et al.* 2002). The encoded uORF from bean reported in this work, was found to be homologous to uORFs from several plants (Fig. 3B).

Differential expression of *Pvadc*, *Pvodc*, *Pvsamdc* and *Pvspd*s in different bean tissues: Transcript levels of *adc*, *odc*, *samdc* and *spds* were analysed by RT-PCR in different bean tissues (Fig. 4). The maximal expression levels of *Pvadc* transcript in non reproductive tissues were observed in young leaves, stems and roots, while no expression was observed in roots tips. We also analysed the transcript levels in flower tissues during the pre-anthesis stage observing that stigma, ovaries and filaments showed the highest expression levels followed by anthers and petals (Fig. 4A). Previously it has been reported that *adc* transcript accumulation appears to be tissue-specific and developmentally-regulated in tomato (Rastogi *et al.* 1993) and soybean (Nam *et al.* 1997). In pea, *adc* gene was highly expressed in young developing tissues such as shoot tips, young leaflets and flower buds, but lower levels were detected in mature and fully expanded leaflets and roots (Perez-Amador *et al.* 1995). A similar behavior was observed in our study where the

highest accumulation of the *adc* transcript occurred in young developmental organs during cell expansion or elongation. ODC activity is essential for normal cell growth and has been closely associated with cellular proliferation.




Fig. 4. Expression analyses of *adc*, *odc*, *samdc* and *spds* in different tissues of *P. vulgaris*. 45 d-old bean plants were used for RT-PCR analyses. 0.01 cm³ RT-PCR product was loaded on each lane and separated by electrophoresis on 1 % agarose gel. Actin product was used as a loading control.

We detected *Pvodc* expression in roots, and surprisingly we observed two weak amplification products in young leaves, suggesting that two *odc* genes could be present in bean (Fig. 4B). Our transcription results correlate with those from *D. stramonium* (Michael *et al.* 1996), where *odc* mRNA accumulates to a greater extent in highly branching and fast growing transformed root cultures as compared to leaves or stems, where cell division is limited. Studies on tomato have shown similar results for *odc* (Kwak and Lee 2001), higher expression levels were detected in roots followed by stems, whereas lower levels were found in young and mature leaves. Alabadí and Carbonell (1998) found that unpollinated ovaries in tomato at 1 d post-anthesis showed a high level of *odc* mRNA, which gradually decreased with time. In contrast, we did not detect *odc* expression in any of the pre-anthesis flower tissues analysed. The *Pvsamdc* transcript was present in rather similar amounts in the majority of the analysed tissues from the bean plant, although ovaries and anthers showed slightly lower levels (Fig. 4C). In potato, *samdc* was highly expressed in young and actively dividing tissues whereas its levels were reduced in mature and non-dividing ones (Mad Arif

et al. 1994). In a study of pea *samdc* (Marco and Carrasco 2002) it was observed that transcript levels varied among tissues and in response to developmental conditions. The levels of expression of *Pvspds* were highest in roots and only barely present in mature and young leaves and stem, whereas no expression was observed in root tips (Fig. 4D). In flower tissues the highest expression was observed in stigma and filaments, followed by ovaries, but no expression was detected in petals and anthers. In contrast, *spds* transcripts were expressed in all plant organs of *A. thaliana* (Hanzawa *et al.* 2002).

In general, we observed the highest accumulation of the analysed *P. vulgaris* transcripts in roots, followed by stigma and filaments, whereas the lowest expression occurred in petals and anthers. These results indicate that expression levels of the genes involved in the biosynthetic pathway of polyamines are tissue-specific and developmentally-regulated. In addition these results support the idea that distribution patterns of genes involved in polyamine biosynthesis do not follow a common pattern in different plant species. This is probably the result of different regulation mechanisms or environmental conditions.

References

Alabadi, D., Carbonell, J.: Expression of ornithine decarboxylase is transiently increased by pollination, 2,4-dichlorophenoxyacetic acid, and gibberellic acid in tomato ovaries. - *Plant Physiol.* **118**: 323-328, 1998.

Ekstrom, J.L., Mathews, I.I., Stanley, B.A., Pegg, A.E., Ealick, S.E.: The crystal structure of human S-adenosylmethionine decarboxylase at 2.25 Å resolution reveals a novel fold. - *Structure Folding Design* **7**: 583-595, 1999.

El Ghachoui, N., Martin-Tanguy, J., Paynot M., Gianinazzi, S.: First report of the inhibition of arbuscular mycorrhizal infection of *Pisum sativum* by specific and irreversible inhibition of polyamine biosynthesis or by gibberellic treatment. - *FEBS Lett.* **385**: 189-192, 1996.

Feirer, R.P., Mignon, G., Litvay, J.D.: Arginine decarboxylase and polyamines required for embryogenesis in the wild carrot. - *Science* **223**: 1433-1435, 1984.

Franceschetti, M., Hanfrey, C., Scaramagli, S., Torrigiani, P., Bagni, N., Burtin, D., Michael, A.J.: Characterization of monocot and dicot plant S-adenosyl-L-methionine decarboxylase gene families including identification in the mRNA of a highly conserved pair of upstream overlapping open reading frames. - *Biochem. J.* **353**: 403-409, 2001.

Galston, A.W., Kaur-Sawhney, R.: Polyamines. - *Plant Physiol.* **94**: 406-410, 1990.

Hanfrey, C., Franceschetti, M., Mayer, M., Illingworth, C., Michael, A.J.: Abrogation of upstream open reading frame-mediated translational control of a plant S-adenosylmethionine decarboxylase results in polyamine disruption and growth perturbations. - *J. biol. Chem.* **277**: 44131-44139, 2002.

Hanzawa, Y., Imai, A., Michael, J.A., Komeda, Y., Takahashi, T.: Characterization of the spermidine synthase-related gene family in *Arabidopsis thaliana*. - *FEBS Lett.* **527**: 176-180, 2002.

Higgins, D.G., Sharp, P.M.: Clustal: a package for performing multiple sequence alignments on a microcomputer. - *Gene* **73**: 237-244, 1988.

Jarvis, B.C., Yasmin, S., Coleman, M.T.: RNA and protein metabolism during adventitious root formation in stem cuttings of *Phaseolus aureus* cv. Berkin. - *Physiol. Plant.* **64**: 53-59, 1985.

Jiménez-Bremont, J.F., Camacho-Villasana, Y.M., Cabrera-Ponce, J.L., Barba de la Rosa, A.P., Ochoa-Alejo, N.: Sequence comparison of plant ornithine decarboxylases reveals high homology and lack of introns. - *Biol. Plant.* **48**: 193-198, 2004.

Kakkar, R.K., Nagar, P.K.: Polyamine and senescence of maintenance foliage of tea [*Camellia sinensis* L.(O) Kuntze]. - *Biol. Plant.* **38**: 153-157, 1996.

Kakkar, R.K., Nagar, P.K.: Distribution and changes in endogenous polyamines during winter dormancy in tea (*Camellia sinensis* L. (O) Kuntze). - *J. Plant Physiol.* **151**: 63-67, 1997.

Kakkar, R.K., Rai, V.K.: Plant polyamines in flowering and fruit ripening. - *Phytochemistry* **33**: 1281-1288, 1993.

Kakkar, R.K., Nagar, P.K., Ahuja, P.S., Rai, V.K.: Polyamines and plant morphogenesis. - *Biol. Plant.* **43**: 1-11, 2000.

Kumar, A., Altabella, T., Taylor, M.A., Tiburcio, A.F.: Recent advances in polyamine research. - *Trends Plant Sci.* **2**: 124-130, 1997.

Kwak, S., Lee, S.H.: The regulation of ornithine decarboxylase gene expression by sucrose and small upstream open reading frame in tomato (*Lycopersicon esculentum* Mill.). - *Plant Cell Physiol.* **42**: 314-323, 2001.

Lee, Y.S., Cho, Y.D.: Identification of essential active-site residues in ornithine decarboxylase of *Nicotiana glutinosa* decarboxylating both L-ornithine and L-lysine. - *Biochem. J.* **360**: 657-665, 2001.

Mad Arif, S.A., Taylor, M.A., George, L.A., Butler, A.R., Burch, L.R., Davies, H.V., Stark, M.J.R., Kumar, A.:

Characterization of the S-adenosylmethionine decarboxylase (SAMDC) gene of potato. - *Plant mol. Biol.* **26**: 327-338, 1994.

Malmberg, R.L., Watson, M.B., Galloway, G.L., Yu, W.: Molecular genetic analysis of plant polyamines. - *Crit. Rev. Plant Sci.* **17**: 199-224, 1998.

Marco, F., Carrasco, P.: Expression of the pea S-adenosylmethionine decarboxylase gene is involved in developmental and environmental responses. - *Planta* **214**: 641-647, 2002.

Martin-Tanguy, J.: Conjugated polyamines and reproductive development: biochemical, molecular and physiological approaches. - *Physiol. Plant.* **100**: 675-688, 1997.

Michael, A.J., Furze, J.M., Rhodes, M.J.C., Burtin, D.: Molecular cloning and functional identification of a plant ornithine decarboxylase cDNA. - *Biochem. J.* **314**: 241-248, 1996.

Moore, R.C., Boyle, S.M.: Nucleotide sequence and analysis of the *speA* gene encoding biosynthetic arginine decarboxylase in *Escherichia coli*. - *J. Bacteriol.* **172**: 4631-40, 1990.

Muchitch, M.J., Edwards, L.A., Fletcher, J.S.: Influence of diamines and polyamines on the senescence of plant suspension cultures. - *Plant Cell Rep.* **2**: 82-84, 1983.

Nam, K.H., Lee, S.H., Lee, J.H.: Differential expression of ADC mRNA during development and upon acid stress in soybean (*Glycine max*) hypocotyls. - *Plant Cell Physiol.* **38**: 1156-1166, 1997.

O'Gara, M., McCloy, K., Malone, T., Cheng, X.: Structure-based sequence alignment of three AdoMet-dependent DNA methyltransferases. - *Gene* **157**: 135-138, 1995.

Perez-Amador, M.A., Carbonell, J., Granell, A.: Expression of arginine decarboxylase is induced during early fruit development and in young tissues of *Pisum sativum* (L.). - *Plant mol. Biol.* **28**: 997-1009, 1995.

Rastogi, R., Dulson, J., Rothstein, S.J.: Cloning of tomato (*Lycopersicon esculentum* Mill.) arginine decarboxylase gene and its expression during fruit ripening. - *Plant Physiol.* **103**: 829-834, 1993.

Rechsteiner, M., Rogers, S.W.: PEST sequences and regulation by proteolysis. - *Trends Biochem. Sci.* **21**: 267-271, 1996.

Sandmeier, E., Hale, T.I., Christen, P.: Multiple evolutionary origin of pyridoxal-5'-phosphate-dependent amino acid decarboxylases. - *Eur. J. Biochem.* **221**: 997-1002, 1994.

Sekowska, A., Bertin, P., Danchin, A.: Characterization of polyamine synthesis pathway in *Bacillus subtilis* 168. - *Mol. Microbiol.* **29**: 851-858, 1998.

Tabor, C.W., Tabor, H.: Polyamines. - *Annu. Rev. Biochem.* **53**: 749-790, 1984.

Tiburcio, A.F., Kaur-Sawhney, R., Galston, A.W.: Polyamine metabolism. - In: Miflin, B.J., Lea, P.J., (ed.): *The Biochemistry of Plants, Intermediary Nitrogen Fixation*. Pp. 283-325. Academic Press, New York 1990.

Watson, M.B., Emory, K.K., Piatak, R.M., Malmberg, R.L.: Arginine decarboxylase (polyamine synthesis) mutants of *Arabidopsis thaliana* exhibit altered root growth. - *Plant J.* **13**: 231-239, 1998.

Yoshida, I., Yamagata, H., Hirasawa, E.: Blue- and red-light regulation and circadian control of gene expression of S-adenosylmethionine decarboxylase in *Pharbitis nil*. - *J. exp. Bot.* **50**: 319-326, 1999.