

BRIEF COMMUNICATION

Lanthanum relieves salinity-induced oxidative stress in *Saussurea involucrata*C.-M. XU^{1,2}, B. ZHAO¹, X.-D. WANG¹ and Y.-C. WANG^{1*}*Institute of Process Engineering, Chinese Academy of Sciences, Beijing 100080, P.R. China¹
Graduate School of Chinese Academy of Sciences, Beijing 100049, P.R. China²***Abstract**

Changes in growth, physiological and biochemical characteristics under salt stress with or without La^{3+} treatment in *Saussurea involucrata* Kar. et Kir. were investigated. The results showed that La^{3+} relieved the plant growth inhibition, improved the leaf water potential and water content, increased the soluble protein and the proline contents and decreased malondialdehyde content under salt stress. Further, addition of La^{3+} significantly increased the activities of superoxide dismutase, ascorbate peroxidase, catalase, and glutathione reductase, decreased the photosynthetic pigment decomposition and increased the ratio of total chlorophyll to carotenoids under salt stress.

Additional key words: antioxidative enzymes, carotenoids, chlorophyll, growth, leaf water potential, proline, proteins.

Saussurea involucrata Kar. et Kir. is an endangered medicinal plant growing in the Tianshan and Kunlun mountain ranges of China, and is very sensitive to salinity stress. High salinity causes both water stress and ionic toxicity, which disputes the various enzymes, function of photosynthetic apparatus and many other processes and severely affects yield and quality of this medicinal plant (Chen *et al.* 1999). It is well known that rare-earth elements form a special group and have interesting effects on plant growth (He and Loh 2000, Ouyang *et al.* 2003). Lanthanum is a rare-earth element and can relieve oxidative damage in plants exposed to a variety of environmental stresses (Gao and Chen 2005, Shi and Chen 2005, Wei *et al.* 1997). Feng *et al.* (1999) reported that La^{3+} could alleviate the decrease of osmotic potential, increase the content of proline and reduce the accumulation of MDA under osmotic stress in the leaves of maize seedlings. There is salt stress affecting the growth of *S. involucrata* in saline area. Up to now, there is few reports on the role of La^{3+} in alleviating the oxidative damage in this plant under salt stress. So, study on this may improve the endangered plant growth and production. In the present study, the effect of La^{3+} on

oxidative damage in *S. involucrata* under salt stress was investigated. The investigation also goes in some way towards studying the effects of La^{3+} on plant growth, soluble protein content and photosynthetic pigments in this species in response to salinity stress.

Seeds of *Saussurea involucrata* Kar. et Kir. were surface-disinfested and aseptically planted into *Vermiculite*, then they were irrigated with liquid NaCl solution at different concentrations (0, 20, 50 and 100 mM) on 0, 5th, 10th and 15th day. The 15-d-old germinated seedlings were divided into two groups. Previous reports indicated that 0.1 mM La^{3+} was more suitable concentration for alleviating environment stress (Pang *et al.* 2002, Zhang *et al.* 2003). So, in this experiment, treatment group seedlings under varying salinity stress were sprayed with 100 cm³ 0.1 mM $\text{La}(\text{NO}_3)_3 \cdot 6 \text{H}_2\text{O}$ solution once a day for five consecutive days. The control group seedlings under varying salinity stress were sprayed with the same volume water. The plants were cultivated under natural light in a greenhouse at 25 °C. At the end of the treatments, plants were harvested, rinsed three times in sterile distilled water, and divided into shoots and roots for further analyse and determination.

Received 10 March 2006, accepted 25 July 2006.

Abbreviations: APOX - ascorbate peroxidase; CAT - catalase; d.m. - dry mass; f.m. - fresh mass; EDTA - ethylenediaminetetraacetic acid; GR - glutathione reductase; MDA - malondialdehyde; NADP - nicotinamide adenine dinucleotide phosphate; ROS - reactive oxygen species; SOD - superoxide dismutase; WC - water content, ψ_w - leaf water potential.

Acknowledgement: This research is supported by Directional Project of Knowledge Innovation Engineering CAS (KGCX2-SW-506).

* Corresponding author; fax: (+86) 10 82627059, e-mail: xocom1978@sohu.com

Length and fresh mass (f.m.) of the shoots and roots were recorded. Then fresh materials were incubated in an oven at 60 °C for 48 h and dry mass (d.m.) was recorded. Water content (WC) was calculated as:

$$WC = (f.m. - d.m.) / f.m. \times 100.$$

Five leaves from each treatment were sampled to measure leaf water potential (ψ_w) using plant pressure chamber (PMS Instrument Company, Model 1000, Albany, USA) method (Gong *et al.* 2003).

Fresh shoots of seedlings were used for biochemical analyses. Soluble proteins were estimated by the method of Bradford (1976) using bovine serum albumin as a standard. The contents of photosynthetic pigments were determined according to Lichtenthaler and Wellburn (1983) in 80 % acetone extract. Proline was estimated according to Bates *et al.* (1973). Lipid peroxidation was determined by measuring malondialdehyde (MDA) formation using the thiobarbituric acid method of Ohkawa *et al.* (1979).

For enzyme assay, fresh shoots (0.2 g) were homogenized in 10 cm³ chilled extraction buffer consisting of 0.1 M phosphate buffer, pH 7.5, 0.5 mM EDTA for superoxide dismutase (SOD; EC 1.15.1.1) and catalase (CAT; EC 1.11.1.6), or 0.1 M phosphate buffer, pH 7.5, 0.5 mM EDTA, 1 mM ascorbic acid in case of ascorbate peroxidase (APOX; EC 1.11.1.11). The homogenate was centrifuged at 15 000 *g* for 20 min and the supernatant fraction was used for the assays. All operations were carried out at 4 °C. All spectrophotometric analyses were performed with a SP-2000 UV spectrophotometer (Shanghai Precision and Scientific Instrument Co., Shanghai, China) at room temperature. SOD activity was estimated at 560 nm according to the method of Beauchamp and Fridovich (1971). APOX activity was determined by measuring the decrease in absorbance of the oxidized ascorbate at 290 nm, according to Nakano and Asada (1981). CAT was determined by measuring the decrease in absorbance of hydrogen peroxide at 240 nm (Aebi 1984). Glutathione reductase (GR; EC 1.6.4.2) activity was determined based on the decrease in absorbance at 340 nm due to the oxidation of NADPH to NADP (Fryer *et al.* 1998).

All the experiments were repeated three times. In each treatment, at least five replicates were used. Prior to statistical analyses, the percentage data were transformed into arc sine square roots. The statistical analysis was performed using the PROC ANOVA of SAS version 6.12.

Salt treatment significantly retarded the growth of seedlings. The length of shoots and roots was remarkably reduced under different salt concentrations. In the case of 100 mM NaCl, the length of shoots and roots was 23.3 and 23.1 % of the lengths without NaCl treatment respectively. Lanthanum application considerably increased the length of shoots and roots under salt stress (Table 1). The mean f.m. and d.m. of a single shoot and root decreased under different salt treatment. La³⁺ application markedly alleviated the plant growth inhibition under salt stress (Table 1). The effects of salinity on ψ_w varied with the salt concentration to which

seedlings exposed. Salt treatment, when the salt concentrations were in the range from 0 to 100 mM, decreased the ψ_w significantly (from -0.98 to -1.53 MPa), whereas La³⁺ application alleviated this trend (Table 1). Without La³⁺ treatment, the leaf WC was considerably reduced from 83.6 % at 0 mM NaCl treatment to 74.6 % at 100 mM NaCl treatment. La³⁺ application alleviated the effects of salt on water content, which reduced from 87.0 to 81.3 % (Table 1). Similar results were observed in the study of lanthanum effects on plantlet differentiation and growth of *Prunus pseudocerasus* Lindl (Li *et al.* 2005).

Soluble protein content declined significantly under salt stress. However, the decline in La³⁺ treated group was lower than that in non-treated group. Under salt stress, proline content significantly increased in many plants (Agarwal and Pandey 2004). The increase of proline with increasing NaCl concentration indicated that the seedlings had self-protective function from the damage by salt stress. At the same NaCl concentration, the addition of La³⁺ promoted proline formation, which indicated that La³⁺ can decrease the damage of salt stress. Lipid peroxidation often acted as an indicator of oxidative damage under salt stress and was determined by measuring MDA content, which was related to lipid membrane damage (Bandoğlu *et al.* 2004). Compared to the control, La³⁺ application alleviated the increasing trend of MDA content (Table 1).

In reactive oxygen species (ROS) scavenging system, the primary scavenger is SOD. SOD activities increased under different NaCl concentrations, but at the same NaCl concentration, La³⁺ application significantly increased the SOD activities, especially at 50 mM NaCl. In the ascorbate-glutathione cycle, APOX reduces H₂O₂ using ascorbate as an electron donor. APOX activities increased significantly with increasing NaCl concentration. The results of the present study were in conformity with those obtained by Sairam *et al.* (2006). Compared with non-lanthanum treatment, APOX activities increased 60.9 and 52.7 % at 50 mM and 100 mM NaCl with La³⁺ treatment, respectively. CAT is another enzyme which is also involved in the detoxification of H₂O₂ by converting H₂O₂ into water and oxygen (Mandhania *et al.* 2006). CAT activities increased under salt stress. Moreover, there was significant difference between with and without La³⁺ application. Compared with non-lanthanum treatment group, GR activities increased 57.9 and 51.6 % at 50 mM and 100 mM NaCl with La³⁺ treatment, respectively (Table 1). Similar results were obtained in different plants such as cucumber (Zhu *et al.* 2004), pea (Hernández *et al.* 2000) and cotton (Gossett *et al.* 1994). So, addition of La³⁺ promoted the activities of the four enzymes, especially at 50 and 100 mM NaCl, increase the antioxidant defence abilities, and alleviate the oxidative damage of proteins and lipids of *S. involucrata* under salt stress.

Salinity greatly decreased chlorophyll content (Table 1) similarly as reported *e.g.* Gadallah and Ramadan (1997). Also the ratio of total chlorophyll to carotenoid content was negatively correlated with NaCl concentration. The

ratio was 7.30 and 7.10 in with and without La^{3+} group at 0 mM NaCl, however, the reduction of the ratio were 11.1 and 22.5 % at 100 mM NaCl concentration, respectively (Table 1). The change may be due to the formation of proteolytic enzymes such as chlorophyllase which is responsible for chlorophyll degradation (Sabater and Rodriguez 1978). Compared to the non-lanthanum treatment, La^{3+} application led to higher ratio of total

chlorophyll to carotenoid content and decreased the photosynthetic pigment decomposition.

It can be drawn a conclusion from above results that lanthanum application promoted the growth of *S. involucrata*, alleviated the oxidative damage and protected the photosynthetic pigments from damage under salt stress.

Table 1. Effects of La^{3+} treatment on growth characteristics, water potential, water content, protein content, proline content, antioxidant enzyme activities and pigment content of *Saussurea involucrata* under different NaCl concentrations. Means $\pm$ SE of 5 replicates.

Parameters	La^{3+} [mM]	NaCl [mM]	20	50	100
		0			
Shoot length	0	11.42 $\pm$ 0.55	6.31 $\pm$ 0.35	4.77 $\pm$ 0.77	2.66 $\pm$ 0.51
[cm]	0.1	14.65 $\pm$ 0.42	8.11 $\pm$ 0.39	7.26 $\pm$ 0.31	5.71 $\pm$ 0.17
Root length	0	9.36 $\pm$ 0.24	6.15 $\pm$ 0.32	4.53 $\pm$ 0.34	2.16 $\pm$ 0.21
[cm]	0.1	12.11 $\pm$ 0.42	7.82 $\pm$ 0.27	6.97 $\pm$ 0.26	5.46 $\pm$ 0.32
Shoot f.m.	0	1.33 $\pm$ 0.02	0.98 $\pm$ 0.05	0.79 $\pm$ 0.04	0.69 $\pm$ 0.03
[g]	0.1	1.68 $\pm$ 0.05	1.31 $\pm$ 0.08	1.06 $\pm$ 0.04	0.96 $\pm$ 0.02
Shoot d.m.	0	0.19 $\pm$ 0.02	0.17 $\pm$ 0.03	0.14 $\pm$ 0.01	0.11 $\pm$ 0.01
[g]	0.1	0.23 $\pm$ 0.03	0.20 $\pm$ 0.02	0.17 $\pm$ 0.01	0.15 $\pm$ 0.01
Root f.m.	0	0.48 $\pm$ 0.03	0.35 $\pm$ 0.01	0.31 $\pm$ 0.02	0.24 $\pm$ 0.01
[g]	0.1	0.57 $\pm$ 0.03	0.44 $\pm$ 0.03	0.35 $\pm$ 0.01	0.33 $\pm$ 0.02
Root d.m.	0	0.05 $\pm$ 0.002	0.04 $\pm$ 0.001	0.04 $\pm$ 0.001	0.03 $\pm$ 0.002
[g]	0.1	0.06 $\pm$ 0.004	0.06 $\pm$ 0.003	0.05 $\pm$ 0.001	0.05 $\pm$ 0.002
ψ_w	0	0.98 $\pm$ 0.05	1.22 $\pm$ 0.05	1.37 $\pm$ 0.07	1.53 $\pm$ 0.05
[-MPa]	0.1	0.87 $\pm$ 0.05	1.06 $\pm$ 0.05	1.18 $\pm$ 0.06	1.35 $\pm$ 0.03
WC	0	83.60 $\pm$ 3.50	81.60 $\pm$ 2.90	76.40 $\pm$ 2.80	74.60 $\pm$ 3.30
[%]	0.1	87.00 $\pm$ 3.50	84.20 $\pm$ 0.40	81.70 $\pm$ 2.90	81.30 $\pm$ 3.50
Protein	0	0.97 $\pm$ 0.03	0.76 $\pm$ 0.01	0.73 $\pm$ 0.02	0.64 $\pm$ 0.04
[mg g ⁻¹ (d.m.)]	0.1	1.13 $\pm$ 0.04	0.97 $\pm$ 0.02	0.90 $\pm$ 0.03	0.77 $\pm$ 0.03
Proline	0	3.17 $\pm$ 0.21	3.98 $\pm$ 0.12	4.41 $\pm$ 0.17	5.48 $\pm$ 0.13
[nmol g ⁻¹ (d.m.)]	0.1	4.67 $\pm$ 0.22	5.36 $\pm$ 0.20	6.02 $\pm$ 0.26	7.93 $\pm$ 0.19
MDA	0	70.40 $\pm$ 5.18	112.68 $\pm$ 4.73	134.17 $\pm$ 4.01	180.19 $\pm$ 8.00
[nmol g ⁻¹ (d.m.)]	0.1	67.52 $\pm$ 3.86	103.21 $\pm$ 4.57	111.46 $\pm$ 5.57	123.57 $\pm$ 3.99
SOD	0	104.90 $\pm$ 5.18	127.30 $\pm$ 6.07	179.50 $\pm$ 7.38	315.60 $\pm$ 8.88
[U g ⁻¹ (d.m.)]	0.1	158.00 $\pm$ 10.16	212.40 $\pm$ 11.84	344.00 $\pm$ 9.61	410.70 $\pm$ 10.42
APOX	0	2.78 $\pm$ 0.84	5.45 $\pm$ 0.76	8.89 $\pm$ 0.60	12.25 $\pm$ 0.67
[U g ⁻¹ (d.m.)]	0.1	5.11 $\pm$ 0.69	10.37 $\pm$ 0.97	14.30 $\pm$ 1.10	18.70 $\pm$ 1.05
CAT	0	314.70 $\pm$ 10.35	331.60 $\pm$ 12.14	396.70 $\pm$ 14.75	481.30 $\pm$ 16.76
[pmol g ⁻¹ (d.m.)]	0.1	507.50 $\pm$ 14.54	535.90 $\pm$ 13.76	553.50 $\pm$ 13.52	719.00 $\pm$ 15.85
GR	0	297.50 $\pm$ 15.25	322.90 $\pm$ 14.25	364.80 $\pm$ 21.96	482.00 $\pm$ 20.60
[U g ⁻¹ (d.m.)]	0.1	472.60 $\pm$ 16.40	464.50 $\pm$ 18.86	575.90 $\pm$ 15.63	730.80 $\pm$ 22.22
Chlorophylls	0	9.02 $\pm$ 0.62	5.82 $\pm$ 0.47	5.12 $\pm$ 0.51	4.92 $\pm$ 0.37
[mg g ⁻¹ (d.m.)]	0.1	11.98 $\pm$ 0.82	8.43 $\pm$ 0.36	7.34 $\pm$ 0.45	6.89 $\pm$ 0.43
Carotenoids	0	1.27 $\pm$ 0.05	0.96 $\pm$ 0.06	0.91 $\pm$ 0.05	0.89 $\pm$ 0.38
[mg g ⁻¹ (d.m.)]	0.1	1.64 $\pm$ 0.04	1.27 $\pm$ 0.04	1.17 $\pm$ 0.05	1.06 $\pm$ 0.04
Chl/Car	0	7.10	6.04	5.61	5.50
	0.1	7.30	6.64	6.29	6.49

References

- Aebi, H.: Catalase *in vitro*. - Methods Enzymol. **105**: 121-126, 1984.
- Agarwal, S., Pandey, V.: Antioxidant enzyme responses to NaCl stress in *Cassia angustifolia*. - Biol. Plant. **48**: 555-

- 560, 2004.
- Bandeoglu, E., Eyidoğan, F., Yücel, M., Öktem, H.A.: Antioxidant responses of shoots and roots of lentil to NaCl salinity stress. - *Plant Growth Regul.* **42**: 69-77, 2004.
- Bates, L.S., Waldren, R.P., Teare, I.D.: Rapid determination of free proline for water-stress studies. - *Plant Soil.* **39**: 205-207, 1973.
- Beauchamp, C., Fridovich, I.: Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. - *Anal. Biochem.* **44**: 276-287, 1971.
- Bradford, M.: A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. - *Anal. Biochem.* **72**: 248-254, 1976.
- Chen, F.J., Yang, Y.G., Zhao, D.X., Gui, Y.L., Guo, Z.C.: Advances in studies of species habitats distribution and chemical composition of snow lotuses (*Saussurea*) in China. - *Chin. Bull. Bot.* **16**: 561-566, 1999.
- Feng, W.X., Zhang, Y.E., Wang, Y.G., Li, F.X.: The alleviative effects of LaCl_3 on the osmotic stress in maize seedlings. - *Henan Sci.* **17**: 45-46, 1999.
- Fryer, M.J., Andrews, J.R., Oxborough, K., Blowers, D.A., Baker, N.R.: Relationship between CO_2 assimilation, photosynthetic electron transport, and active O_2 metabolism in leaves of maize in the field during periods of low temperature. - *Plant Physiol.* **116**: 571-580, 1998.
- Gadallah, M.A.A., Ramadan, T.: Effects of zinc and salinity on growth and anatomical structure of *Carthamus tinctorius* L. - *Biol. Plant.* **39**: 411-418, 1997.
- Gao, Y.S., Chen, J.S.: Effects of La^{3+} on antioxidant system in wheat seedlings leaves under salt stress. - *J. chin. rare Earth Soc.* **23**: 490-495, 2005.
- Gong, H.J., Chen, K.M., Chen, G.C., Zhu, X.Y., Wang, S. M., Zhang C.L.: Effects of gradual drought on the fatty acid composition of polar lipids, H^+ -ATPase and 5'-AMPase activities in the plasma membranes of two spring wheat leaves. - *Acta phytoecol. sin.* **27**: 459-465, 2003.
- Gossett, D.R., Millhollon, E.P., Lucas, M.C.: Antioxidant response to NaCl stress in salt tolerant and salt sensitive cultivars of cotton. - *Crop Sci.* **34**: 706-714, 1994.
- He, Y.W., Loh, C.S.: Cerium and lanthanum promote floral initiation and reproductive growth of *Arabidopsis thaliana*. - *Plant Sci.* **159**: 117-124, 2000.
- Hernández, J.A., Jiménez, A., Mullineaux, P., Sevilla, F.: Tolerance of pea (*Pisum sativum* L.) to long-term salt stress is associated with induction of antioxidant defences. - *Plant Cell Environ.* **23**: 853-862, 2000.
- Li, H.S., Cui, D.H., Wang, J.: Effect of La on plantlet differentiation and growth of *Prunus pseudocerasus* Lindl. - *Plant Nutr. Fert. Sci.* **11**: 140-142, 2005.
- Lichtenthaler, H.K., Wellburn, A.R.: Determinations of total carotenoids and chlorophylls *a* and *b* of leaf extracts in different solvents. - *Biochem. Soc. Trans.* **11**: 591-592, 1983.
- Mandhanian, S., Madan, S., Sawhney, V.: Antioxidant defense mechanism under salt stress in wheat seedlings. - *Biol. Plant.* **50**: 227-231, 2006.
- Nakano, Y., Asada, K.: Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. - *Plant Cell Physiol.* **22**: 867-880, 1981.
- Ohkawa, H., Ohishi, N., Yagi, Y.: Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. - *Anal. Biochem.* **95**: 351-358, 1979.
- Ouyang, J., Wang, X.D., Zhao, B., Yuan, X.F., Wang, Y.C.: Effects of rare-earth elements on the growth of *Cistanche deserticola* cells and the production of phenylethanoid glycosides. - *J. Biotechnol.* **102**: 129-134, 2003.
- Pang, X., Wang, D.H., Xing, X.Y., Peng, A., Zhang, F.S., Li, C.J.: Effect of La^{3+} on the activities of antioxidant enzymes in wheat seedlings under lead stress in solution culture. - *Chemosphere* **47**: 1033-1039, 2002.
- Sabater, B., Rodriguez, M.T.: Control of chlorophyll degradation in detached leaves of barley and oat through effect of kinetin on chlorophyllase levels. - *Physiol. Plant.* **43**: 274-276, 1978.
- Sairam, R.K., Srivastava, G.C., Agarwal, S., Meena, R.C.: Differences in antioxidant activity in response to salinity stress in tolerant and susceptible wheat genotypes. - *Biol. Plant.* **49**: 85-91, 2005.
- Shi, P., Chen, G.C.: Effects of La^{3+} on the active oxygen-scavenging enzyme activities in cucumbers seedling leaves. - *Russ. J. Plant Physiol.* **52**: 294-297, 2005.
- Wei, A.L., Chen, Y.Z., Wang, Y.G.: Influences of $\text{La}(\text{NO}_3)_3$ on alleviating NaCl induced injury effects in soybean seedling and its physiological and biochemical reactions. - *J. Shanxi agr. Univ.* **17**: 103-108, 1997.
- Zhang, L., Zeng, F., Xiao, R.: Effect of lanthanum ions (La^{3+}) on the reactive oxygen species scavenging enzymes in wheat leaves. - *Biol. Trace Element Res.* **91**: 243-245, 2003.
- Zhu, Z.J., Wei, G.Q., Li, J., Qian, Q.Q., Yu, J.Q.: Silicon alleviates salt stress and increases antioxidant enzymes activity in leaves of salt-stressed cucumber (*Cucumis sativus* L.). - *Plant Sci.* **167**: 527-533, 2004.