

REVIEW

Chloroplast-specific universal primers and their uses in plant studies

N. HAIDER*

*Department of Molecular Biology and Biotechnology, Atomic Energy Commission of Syria,
P.O. Box 6091, Damascus, Syria***Abstract**

Universal (consensus) primers are those primers that have the ability to amplify the targeted region of DNA across a broad range of individuals in a certain group of organisms. In plants, such universal primers have been designed to target regions in the nuclear, mitochondrial or chloroplast genome. Among these three genomes, the chloroplast genome is the most suited for the design of consensus primers due to the lower rate of evolution and hence conservation of gene order and sequence of the genome among the different plant species compared to the other two genomes. Several molecular studies in plants have developed and used chloroplast-specific universal primers. In this review, I present some examples of the nuclear DNA-specific universal primers and discuss the features of the chloroplast DNA that make it the most suited for the design of such primers. I then refer to all chloroplast-specific primers developed so far and provide some examples of molecular studies and applications that made use of them.

Additional key words: chloroplast genome, consensus primers, molecular systematics.

Introduction

Several molecular systematic studies have made use of primers that are described as being 'universal'. In such instances, the term 'universal' was generally used to describe the ability of those primers to amplify a target DNA sequence efficiently across a range of individuals belonging to a particular group of organisms (Haider 2003). The taxonomic rank at which the primers apply varies with different sets of primers reported to have

different levels of universality (taxonomic spread). For example, primers designed by Taberlet *et al.* (1991) are apparently able to amplify products using template DNA from species across the whole plant kingdom. Such primers have proved valuable for molecular systematic studies since they save time, effort and cost that would otherwise be required to identify and sequence a suitable locus across all target organisms.

Nuclear DNA-based universal primers

The internal transcribed sequence (ITS) forms a part of the nuclear genome that contains multiple copies of coding regions for rDNA genes that are arranged in several tandem arrays (Appels and Honeycutt 1986). The ITS region has been commonly used as a target for the design and use of universal primers in plant research. Indeed, there is now a set of seven ITS universal primers (White *et al.* 1990, Sun *et al.* 1994) that are widely

applied for diagnostic (Linder *et al.* 2000) and phylogenetic studies in many angiosperm families (Baldwin *et al.* 1995). In 2005, Moore and Field were able to identify the roots of different plant species in mixed samples using sequences of the nuclear ITS region. Sequencing of the same region proved useful for DNA barcoding of members of the *Cycadales* (Sass *et al.* 2007).

Received 11 November 2009, *accepted* 25 May 2010.

Abbreviations: cp - chloroplast; ETS - external transcribed spacer; GM - genetically modified; IGS - intergenic spacer; ITS - internal transcribed sequence, NTS - nontranscribed spacer; SSR - simple sequence repeats or microsatellites; STS - sequence-tagged site.

Acknowledgements: Many thanks to Prof. Ibrahim Othman, the Director General of Atomic Energy Commission of Syria, for his support. I would also like to thank Prof. Mike J. Wilkinson, Director General of University of Aberystwyth, UK, for his endless advice.

* Fax (+963) 11 6112289, e-mail: scientific@aec.org.sy

The intergenic spacer (IGS), which constitutes of a nontranscribed spacer (NTS) and an external transcribed spacer (ETS) between 26S and 18S rDNA genes, of the nuclear genome has been similarly targeted for the design of universal primers, although these primers have been generally less widely used. Baldwin and Markos (1998) designed two universal primers (18S-IGS and 26S-IGS) for the amplification of the entire IGS region. In the same study, an internal primer within the ETS was developed and used with the primer 18S-IGS for universal amplification of the ETS region in the tribe *Heliantheae* (*Compositae*) and phylogeny reconstruction of *Calycadenia* (*Compositae*). Likewise, the NTS between repeated 5S-ribosomal RNA genes has also been used for the construction of universal primers such as those described by Cox *et al.* (1992), who reported the efficiency of four primers they designed when amplifying target DNA from several distantly related mono-

cotyledonous plants.

Taylor *et al.* (2001) designed primers that target single locus coding regions to generate sequence-tagged site (STS) markers in cereals that could also amplify in *Lolium perenne* for use in comparative mapping. The use of such nuclear universal primers, also allows the acquisition of unknown sequence adjacent to known sequences, both in the upstream and downstream directions (Weber *et al.* 1998). Recently, a set of universal primers has been developed by Prado *et al.* (2004) to clone conserved domains of the acetolactate synthase (ALS) gene in monocotyledonous and dicotyledonous plants. More recently, McIntosh *et al.* (2005) designed three pairs of grass-specific universal primers that amplify the nuclear gene GBSS1. These primers proved to have utility in identification of some grass species and may be also useful in the distinction of species from other plant families as the authors revealed.

Cytoplasmic DNA-based universal primers

The cytoplasm of plants contains two genomes, namely the chloroplast and mitochondrial genomes. The chloroplast genome contrasts strikingly with the nuclear genome. The inheritance of chloroplast DNA (cpDNA) is most commonly clonal (through maternal parent in 80 % of angiosperms; Matsushima *et al.* 2008). It is also not influenced by polyploidy, gene duplication and recombination that are widespread features of the nuclear genomes of plants (Tanksley and Pichersky 1988, Harris and Ingram 1991). Therefore, cpDNA varies little among angiosperms in terms of size, structure and gene content (Curtis and Clegg 1984).

The mitochondrial DNA (mtDNA) bears contrasting features to those of the chloroplast genome that influence their utility for the design and application of universal primers. The mtDNA is large in plants, has a slow nucleotides substitution rate, extensive level of intramolecular recombination and in turn, non-conserved order of genes. Consequently, universal primers that target sequences of mitochondrial loci are more difficult to design and generate fewer polymorphisms. There are therefore only a very limited number of mtDNA-based universal primers that have been developed (Demesure *et al.* 1995, Dumolin-Lapegue *et al.* 1997a). In contrast, the cpDNA features stated above make it ideal for the design of the majority of universal primers available for the molecular analysis of plants. The increasing number of published sequences of the entire chloroplast genome from several diverse plant species (reached 122 in 2010, http://megasun.bch.umontreal.ca/ogmp/projects/other/cp_list.html) that are freely available in DNA sequence databases facilitated the design of such primers. Amplification of target DNA using these primers is usually followed by direct sequencing (*e.g.* Meudt and Bayly 2008) or electrophoresis of either entire or

restricted amplified fragments (*e.g.* Gillespie and Boles 2001).

Universal primers that target the chloroplast simple sequence repeats (cpSSRs): The slower rate of mutation in the chloroplast genome compared to the nuclear genome means that the primers that target the mononucleotide cpSSRs are usually more universal and can often amplify across related species (Provan *et al.* 2001). For instance, Vendramin *et al.* (1996) designed universal primers for the amplification of 20 SSRs (microsatellites) in *Pinaceae*. Similarly, five out of seven cpSSR primer pairs that were designed using the complete chloroplast genome of *Oryza sativa* amplified efficiently across other wild and cultivated rice species (Provan *et al.* 1997). Weising and Gardner (1999) likewise designed a set of ten pairs of universal primers to detect mononucleotide repeat variation in cpDNA of dicotyledonous angiosperms, and used the resultant amplicons to reveal intraspecific and interspecific polymorphism within the genera *Nicotiana*, *Lycopersicon* (both *Solanaceae*) and *Actinidia* (*Actinidiaceae*). Such universal primers that can amplify across a broad range of plant families have been also reported. For instance, Cato and Richardson (1996) designed five pairs of cpSSR primers that amplified successfully in eight families of higher plants. In 2003, Chung and Staub designed 23 pairs of cpSSRs consensus primers that amplify targeted regions in a diverse array of plant species. Recently, Provan *et al.* (2004a) developed five pairs of universal primers for the amplification of cpSSRs in grasses (*Poaceae*). More recently, Cheng *et al.* (2006) developed a set of universal cpSSR primers to explore cpDNA diversity among sub-tropical and tropical fruit crops, where cpDNA sequences are unknown.

Universal primers that target the chloroplast ITS (cpITS): The ITS region of chloroplast rDNA have been similarly targeted for development of universal primers. For example, Goremykin *et al.* (1996) designed two primers to amplify the region spanning from the 3' terminus of the 23S rRNA gene to the 5' terminus of the 5S rRNA gene (located in chloroplast inverted repeats (IRs)). These primers amplified successfully the targeted DNA region that includes the 4.5S rRNA gene and the two chloroplast ITS regions (cpITS2 and cpITS3) in 20 angiosperms, seven gymnosperms and 16 ferns. Samigullin *et al.* (1998) showed that these primers as well as the primer pair designed by the authors to target the chloroplast ITS4 (chloroplast 5S rRNA-tRNA (ACG) spacer) to amplify efficiently across 26 bryophytes.

Universal primers that target the chloroplast noncoding regions: Some chloroplast-encoded genes are interrupted by introns (Clegg 1993). These introns and other noncoding cpDNA regions (intergenic spacers) are variable but separate highly conserved regions. Their variability can be exploited in systematic studies in plants even at lower taxonomic levels and so such regions have been targeted for the design of universal primers. The dispersed distribution of transfer RNAs (*trns*) genes over the chloroplast genome, together with their high level of sequence conservation make them ideal targets for the design of universal primers that bind to conserved regions (Petit *et al.* 1996) that flank more variable regions (noncoding regions). Chloroplast-specific universal primers designed so far have various degrees of "universality" varying from family level such as those of Tsumura *et al.* (1996) that amplify 11 different chloroplast genes across *Dipterocarpaceae*, through to those of Taberlet *et al.* (1991) that target noncoding regions of *trnT* (UGU)-*trnL* (UAA) 5'exon (primers a and b), *trnL* (UAA) intron (primers c and d) and *trnL* (UAA) 3'exon-*trnF* (GAA) (primers e and f) and amplify products in the majority of plant species including algae, bryophytes, pteridophytes, gymnosperms and angiosperm. Primers designed by Taberlet *et al.* (1991) to amplify the *trnL* intron have been shown to amplify the targeted intron in many angiosperms (*e.g.* King and Ferris 1998, Horres *et al.* 2000, Zuber and Widmer 2000) and gymnosperm species (*e.g.* Sperisen *et al.* 2000, Terry *et al.* 2000).

Several other universal primers have been reported that amplify noncoding spacers of the chloroplast genome in several angiosperms and gymnosperms. For instance, Fofana *et al.* (1997) designed three pairs of universal primers that target the intergenic spacers of *atpB-rbcL*, *rpsL4-psaB* and *petA-psbE*. Chiang *et al.* (1998) produced universal primers to amplify and sequence a chloroplast noncoding spacer between the *atpB* and *rbcL* genes. The authors suggested that analysis of this spacer would be useful for molecular systematics at the generic, specific, and even subspecific levels to study population genetics

in some plant species. Furthermore, Demesure *et al.* (1995) and Dumolin-Lapegue *et al.* (1997a) designed nine and 16 universal primer pairs, to amplify spacers in the chloroplast and mitochondrial genomes, respectively. In 2001, Grivet *et al.* designed 20 conserved cpDNA primer pairs that target chloroplast intergenic spacers that were tested on 20 plant species belonging to 13 families. Primers developed by Demesure *et al.* (1995) have been proved to amplify in many plant species. For instance, Ziegenhagen and Fladung (1997) were able to PCR-amplify the chloroplast intergenic spacer *trnS-psbC* (Demesure *et al.* 1995) in 62 woody plant species that represented a broad systematic range in both gymnosperms and dicotyledonous angiosperms. Moreover, Sun (2002) indicated that published universal primers by Demesure *et al.* (1995) for amplification of *trnK* (exon 1)-*trnK* (exon 2) successfully amplified the respective regions in *Elymus* species.

Added to those consensus primers mentioned above, Hamilton (1999) developed four primer pairs for the amplification of chloroplast intergenic regions with intraspecific variation. In 2000, Graham and Olmstead developed primers for 14 slowly evolving chloroplast genes that revealed to be efficient in amplifying and sequencing across the seed plants. These primers proved to be a useful tool for inferring the relationships of basal angiosperms and other land plants. In the same year, Nishizawa and Watano (2000) developed universal primer pairs that are suitable for PCR-SSCP analysis of cpDNA in angiosperms. Similarly, Provan *et al.* (2004b) have developed two sets of universal primers for the amplification of chloroplast coding and non-coding regions in green (*Chlorophyta*, eight pairs) and red (*Rhodophyta*, nine pairs) algae. In another study, Lee and Wen (2004) designed universal primers that successfully amplified the entire chloroplast *trnC-trnD* region in different groups of flowering plants. Recently, Watts *et al.* (2008) designed universal primers that amplify and sequence the large Domain IV (D4) loop of the cpDNA in several angiosperm introns. Haider and Wilkinson (in preparation) developed 84 potential universal primers to target 42 regions of the chloroplast genome, and investigated the level of variation in regions flanked by primers designed.

It is worth noting that the design of the striking number of cpDNA-specific universal primers mentioned above was facilitated by the comprehensive DNA sequence information available on the NCBI database (<http://www3.ncbi.nlm.nih.gov/Entrez/nucleotide.html>) for the entire chloroplast genomes of many plant species and more importantly, the availability of nucleotide sequences of chloroplast loci targeted in several plant species representing monocotyledons, dicotyledons and gymnosperms for some loci. When Haider and Wilkinson (in preparation) developed their enlarged set of universal primers, homologous sequences recovered from the NCBI database for each locus targeted by those primers

exhibited, generally, a high level of sequence conservation among plant species. This made the alignment of homologous sequences relatively simple exercise for each

locus, and in turn allowed easy detection of the most highly conserved regions that were selected as primer binding sites.

Uses of chloroplast-specific universal primers in plant molecular studies

The majority of chloroplast universal primers described above and in particular those of Taberlet *et al.* (1991) and Demesure *et al.* (1995) have been used for various taxonomic studies in plants such as:

Evolution and origin of plants: The maternal mode of inheritance of cpDNA is a particular advantage in studying the origin and maternal parent of the hybrid and polyploid species (Palmer 1987). CpDNA, therefore, has been useful for the study of plants origins and evolution. For instance, *Allium fistulosum* was found to have originated monophytically from its close wild relative *A. altaicum* as revealed by RFLP analysis of five noncoding cpDNA regions and RAPD analysis of nuclear DNA (Friesen *et al.* 1999). Fofana *et al.* (1999) also examined an overview on the evolution of the genus *Phaseolus* on the basis of cpDNA diversity using PCR-RFLP. Based on both nuclear DNA and cpDNA data, Friesen *et al.* (2000) unambiguously placed *A. altaicum* within *Allium* subgenus *Rhizirideum*. In another study, comparison of data based on nuclear genome and sequences of the chloroplast regions *trnL*, *trnL-F* and *trnT-L* suggested that *Raphanus sativus* is a hybrid between the *Brassica nigra* and the *B. rapa* × *B. oleracea* lineages with the latter as the maternal parent (Yang *et al.* 2002). Other examples of using universal chloroplast primers to study origin of plants is the study of origin of *Spiranthes diluvialis* (Szalanski *et al.* 2001), *Armeria villosa* ssp. *carratracensis* (Feliner *et al.* 2002), the hybrid *Potamogeton* × *sudermanicus* Hagstr. (Fant *et al.* 2003), some species of kiwifruit (*Actinidia deliciosa*, *Actinidiaceae*) (Chat *et al.* 2004), and the diversity and the evolution of several other legumes (Angioi *et al.* 2009).

Phylogenetics: Molecular analysis offers a useful way to measure the amount of genetic variability in cultivated species and their wild relatives. Evaluation of genetic diversity in turn allows the clarification of phylogenetic relationships among those species and may provide a rationale for choosing strategies for germplasm collection, conservation and use of genetic resources (Buso *et al.* 2001). Most importantly, such information also has value for the improvement of cultivated species, for which the wild relatives are of considerable value because they possess extensive genetic variability. Vakhitov *et al.* (2003) argued that it is important to understand the phylogenetic relationships among cultivated wheat species and their wild relatives for the genetic improvement of these crops.

Palmer (1987) argued that parallelisms and convergences occur so rarely at the species level that it is often possible to build unambiguous phylogenetic using cladistic approaches. Therefore, cpDNA has been widely used for the phylogenetic reconstruction of crop plant species (Olmstead and Palmer 1994). For instance, Lakshmi *et al.* (2002) constructed molecular phylogeny for species of the Indian mangrove tribe *Rhizophoreae* based on sequences of *rbcL*. Chloroplast universal primers available enhance the capacity of evolutionary scientists to determine phylogenetic relationships of both closely or distantly related plant species most commonly using sequence analysis. Noncoding regions for which universal primers are published have also been targeted for phylogenetic studies in 39 species of the tribe *Rubieae* (Natali *et al.* 1995), common beech (*Fagus sylvatica*) (Demesure *et al.* 1996) and eight white oak species (Dumolin-Lapègue *et al.* 1997b). Other examples on using chloroplast universal primers to study phylogenetic relationships are those on the genus *Leonardoxa* (*Leguminosae: Caesalpinioideae*) (Brouat *et al.* 2001), *Phaseolus* (Fofana *et al.* 1997), *Hordeum* (Nishikawa *et al.* 2002), *Hypericum* (Pilepic 2002), *Crinum* (*Amaryllidaceae*) (Meerow *et al.* 2003), *Panax* (Lee and Wen 2004), *Poa* (Gillespie and Soreng 2005), and some *Diospyros* spp. (*Ebenaceae*) including *Diospyros kaki* (Hu *et al.* 2008).

Mode of inheritance: Determination of the mode of the cpDNA inheritance for a group of plant species is useful for studies of species origins, hybridisation, introgression and population genetics (Yang *et al.* 2000). For example, Hollingsworth *et al.* (1999) established the direction of hybridisation and the mode of chloroplast inheritance to investigate spread of invasive weeds belonging to the genus *Fallopia*. Recently, the mode of cpDNA inheritance has been determined in intergeneric hybrids between *Carica papaya* and four different *Vasconcellea* species (Van Droogenbroeck *et al.* 2005).

Feliner *et al.* (2002) similarly provided evidence for a maternal inheritance of cpDNA in *Armeria* based on sequences from two chloroplast regions, namely *trnL-trnF* and *trnD-trnT* that were amplified using published universal primers. In a study carried out by Collins *et al.* (2003), species-specific chloroplast markers developed by analysing restriction digestions of PCR-amplified *trnL-F* fragments confirmed the direction of crosses that generated putative hybrids of three species of *Taxus* (*Taxaceae*), namely *T. baccata*, *T. canadensis* and *T. cuspidate*. Likewise, Panda *et al.* (2003) developed

PCR-RFLP markers that differentiated clearly between sweet cherry (*Prunus avium*) and sour cherry (*P. cerasus*) using six chloroplast universal primer pairs. These markers allowed the authors to understand the maternal inheritance of chloroplast genome in *P. avium* and indicated that the latter is not the maternal ancestral species of *P. cerasus*.

Exclusively uniparental paternal inheritance of cpDNA has been reported to be the commonest in gymnosperms (cited by Rebound and Zeyl 1994) such as red and black spruce (Bobola *et al.* 1996), but is rare amongst angiosperms. Paternal inheritance of the cpDNA has been observed in some angiosperm plant species perhaps slightly compromising the value of analysis of cpDNA. For instance, strict paternal inheritance of cpDNA was found in progeny of interspecific controlled crosses in the genus *Actinidia* (Testolin and Cipriani 1997) based on PCR-RFLP on *trnT-L* and *trnL-F*. Paternal transmission of chloroplast genomes allowed the detection of the paternal genetic lineages of species in the genus. Restriction analysis of three cpDNA regions (*trnT-L*, *trnL-F* and *trnD-T*) that were amplified using universal primers also indicated that the cpDNA inheritance is uniparental and paternal in the genus *Larrea* (*Zygophyllaceae*), which was the fifth genus to have paternal inheritance in angiosperms beside *Medicago*, *Turnera*, *Pharbitis* and *Actinidia* as stated by Yang *et al.* (2000). In a recent study, Matsushima *et al.* (2008) used cpDNA-specific universal primers to determine cpDNA inheritance in *Medicago truncatula*. Based on sequences of the chloroplast *trnL-F*, Zhang *et al.* (2009) were also able to infer the maternal donor of species in the genus *Kengyilia*.

Forensic botany: Compared to the one nucleus within each plant cell, chloroplasts are found in large numbers and each of them is protected by a protein membrane that reduce bacterial breakdown of the cpDNA (Wilkinson and Linacre 2000). The latter believe that this gives the DNA test based on chloroplast genome advantages over tests based on nDNA for forensic purposes. Although, forensic botany can provide significant evidence during criminal investigations, its most common application still, however, limited to identifying specific and suspected illegal plants (Ferri *et al.* 2009).

To date, very few studies, however, have used cpDNA-based evidence in forensic investigations. For instance, nested PCR using universal primers c-d (Taberlet *et al.* 1991) and 'H' primer specific to *Cannabis sativa* (Linacre and Thorpe 1998) allowed Wilkinson and Linacre (2000) to detect the presence of *C. sativa* DNA on the hands of subjects who have recently handled *Cannabis* material and then performed a number of tasks such as rubbing hands on trousers or placing hands in pockets. Bever *et al.* (on-line reference) used DNA sequence variation in *rbcL* to identify the individual plant components found in botanical trace evidence from as

little as a 1 mm punch of dried plant tissue or from five grains of fresh pine pollen. Generally, studies of the identification of species in relation to forensic sciences have used DNA obtained from individual plant specimens of a known species (Linacre and Thorpe 1998). More recently, *rbcL* locus was amplified with universal primers and sequenced across all the elements in a DNA mixture of unknown plant composition (Bever and Cimino 2001). Sequences generated enabled the authors to identify each of the plants as part of the sample. It seems also promising that cpDNA-specific universal primers available may provide a valuable means to generate species-specific markers that can be used in crime investigations such as murder, distribution of drugs, kidnapping, acts of terrorism and sexual assault (Bever *et al.* on-line reference). For example, Wesselink and Kuiper (2008) made a successful attempt for species identification of botanical trace evidence using molecular markers that were generated using cpDNA-specific universal primers. Similarly, Ferri *et al.* (2009) investigated the forensic use of two non-coding chloroplast regions, *psbA-trnH* and *trnL-trnF*, to create a multimarker system for species identification that could be useful throughout the plant kingdom.

Seed dispersal: Seed dispersal has a profound effect on vegetation structure and its dynamics, colonization of new habitats and maintenance of diversity (Wang and Smith 2002). The authors believe that the analysis of molecular markers allows matching dispersed seeds and seedlings with parent plants for an easier detection of long distance dispersal events.

Since cpDNA is maternally transmitted in the majority of angiosperms and thus usually reflects only seed dispersal, using chloroplast markers specific to plant species can therefore provide useful information for the detection of seed dispersal of a plant species to the cultivation area of another species. For example, using universal primers for cpDNA analysis, Petit *et al.* (1997) observed interspecific sharing of the maternal genome of two oak species: *Quercus petraea* and *Q. robur* that strongly suggests that long-distance seed dispersal events followed by interspecific exchanges were involved at the time of colonization of oak species in Europe. In another study, Potgieter and Albert (2001) analysed sequences of the *trnL* intron and *trnL-F* spacer in *Apocynaceae* to resolve phylogenetic relationships in the family that were used as a basis to investigate trends in seed dispersal in *Apocynaceae*. In the same year, the spatial distribution of maternally inherited cpDNA markers over the French part of the range of *Sorbus torminalis* was carried out by Oddou-Muratorio *et al.* (2001) to quantify the relative importance of seed and pollen dispersal. Similarly, using chloroplast universal primers for cleaved amplified polymorphic sequence (CAPS), the spatial mixing of haplotypes in *Quercus robur*, *Q. petraea* and *Q. pubescens* throughout the Swiss Alps and adjacent

regions as a consequence of seed dispersal was low (Matyas and Sperisen 2001). In a recent study, Griffin and Barrett (2004) indicated that seed dispersal between populations of *Trillium grandiflorum* is less likely than pollen flow. Oliver *et al.* (2006) studied seed dispersal of the arctic-montane species *Saxifraga hirculus* (*Saxifragaceae*).

Interspecific variation: Polymorphism that may be detected among different plant species using chloroplast markers can be exploited for the study of interspecific variation that can describe the history of these species. Published chloroplast universal primers have proved useful for the detection of interspecific variation among several plant groups. For instance, PCR-RFLP using universal primers allowed the detection of interspecific variation in various plant genera such as *Leguminosae* (Fofana *et al.* 1997), *Abies* (Parducci and Szmidt 1999) and *Elymus* (Sun 2002). Using universal chloroplast primers referred to as AS, CD and DT (Demesure *et al.* 1995) for PCR and then digestion of amplification products, Pilepic (2002) revealed considerable degree of interspecific polymorphism between *Hypericum perforatum* and *H. maculatum*. In 2004, interspecific differences in the genus *Citrullus* were resolved by Dane *et al.* Genetic variation and relationships among *Ulex* (*Fabaceae*) species in Southern Spain and Northern Morocco were similarly assessed by cpSSR markers (Cubas *et al.* 2005). Using three universal primer pairs, Bouhadida (2007) analyzed cpDNA diversity and revealed genetic relationships in the genus *Prunus*. Recently, Basha and Sujatha (2009) investigated interspecific variation among *Jatropha* species using consensus cpSSR markers.

Somatic hybridisation: Sexual barriers (ploidy and/or endosperm balance number) that prevent interspecific crosses between cultivated species and their wild relatives may restrict exploitation of wild germplasm in breeding programs. Somatic hybridization (fusion of somatic cells in culture) has made many wild species accessible for gene transfer to their cultivated relatives (Bastia *et al.* 2001). The inheritance of chloroplasts in somatic hybrids is uncertain and can vary between experiments or species combinations. Chloroplast-specific universal primers can be used as an easy tool to follow the fate of chloroplasts after somatic fusion of cells from two species. Bastia *et al.* (2000) developed somatic hybrids *via* protoplast fusion between *Solanum tuberosum* and *S. commersonii* and used cpRFLP to characterize chloroplast transmission into the hybrids. Similarly, Bastia *et al.* (2001) used universal primers to amplify intronic and intergenic regions to determine the chloroplast genome composition of interspecific somatic hybrids in the genera *Solanum* and *Brassica*. Cheng *et al.* (2003) also analysed cpDNA inheritance of 10-year-old intergeneric somatic hybrids trees between *Citrus sinensis* and *Fortunella crassifolia*

using CAPS that implied universal primers. In another study, PCR-RFLP using universal primers has been applied for the analysis of the cytoplasmic constitution of *Aurantioideae* intergeneric somatic hybrids (Lotfy *et al.* 2003). Fu *et al.* (2004) analysed the cytoplasmic genomes of intergeneric somatic hybrid combining Goutou sour orange and *Poncirus trifoliata* using universal primers. Recently, Olivares-Fuster *et al.* (2007) used cpDNA universal primers to characterize somatic hybrids and cybrids obtained by fusion of *Citrus sinensis* and *C. excelsa* protoplasts.

Conservation: The generation of species-specific markers using chloroplast universal primers may be useful for studies of the maintenance of germplasm collections. Such repositories are valuable for the conservation and use of plant genetic resources for the improvement of agricultural and forestry crops. In this context, the primers can be used to: 1) ensure identity and integrity of accessions held within a germplasm collection in a genebank, 2) characterisation of seeds entering a genebank, 3) checking seeds held in genebanks if they are contaminated with seed of other species and 4) detect any introgressive hybridization in *ex situ* collections. They can be also exploited to protect wild species of crop species in their natural habitats (*in situ* conservation). For example, sequencing of the chloroplast intron *trnL* and the intergenic spacer *trnL-F* were used to investigate the genetic variability in *Milicia* from three West African countries to design conservation strategies for species in this genus (Ofori *et al.* 2001). In 2004, Honjo *et al.* carried out a study in Japan on the intraspecific phylogeography of the endangered species *Primula sieboldii* to help design conservation programs. Similarly, Aoki *et al.* (2006) illustrated how sequencing of noncoding regions of cpDNA that were amplified using universal primers for phytogeographic studies could provide a basis for identifying suitable units for conservation. Recently, using cpDNA-specific universal primers, De Lange *et al.* (2008) carried out a case study with two enigmatic and uncommon species of *Crassula* from New Zealand for biosystematics and conservation purposes.

Spontaneous hybridisation: Hybridization is an important evolutionary phenomenon, and therefore a better understanding of the dynamics of interspecific gene flow and resulting morphological and genetic patterns has been of widespread interest (Luhová *et al.* 2007).

Interspecific hybridisation and introgression have been reported to be common phenomena in many plant species, *e.g.* *Chaenomeles cathayensis* and *C. speciosa* (Bartish *et al.* 2000) or even genera, *e.g.* *Brassica napus* and *Raphanus raphanistrum* (Eber *et al.* 1994). Hence, natural hybridisation can occur between cultivated plants and their wild relatives when they come into contact (Ellstrand *et al.* 1999).

Numerous studies have attempted to characterise or quantify the occurrence of spontaneous hybrids in natural habitats (e.g. Scott and Wilkinson 1998). In works of this kind, it is clearly important to be able to distinguish both parental species from the hybrid and also from each other. Bartish *et al.* (2000) was able to detect spontaneous hybridisation between *Chaenomeles cathayensis* and *C. speciosa*. Interest in the quantification of interspecific hybridisation has increased dramatically with the commercial cultivation of genetically modified (GM) crops, where attention is focussed on the ecological consequences of transgene recruitment by wild relatives of GM crops by gene flow (Lavigne *et al.* 2002).

Amplification and sequencing of the *trnL-trnF* region (Taberlet *et al.* 1991) of cpDNA provided an evidence of hybridization between two polyploid *Cardamine* (*Brassicaceae*) species in North-western Spain (Luhová *et al.* 2007). Based on sequence information generated from the same chloroplast region, a natural hybrid between *Ligularia paradoxa* and *L. duciformis* (*Asteraceae*, *Senecioneae*) from Yunnan, China was recovered (Pan *et al.* 2008). In 2009, Csencsics *et al.* carried out a large scale survey of *Populus nigra* presence and genetic introgression from non-native poplars in Switzerland based on molecular identification using universal primers for the analysis of cpDNA. Similarly, Haider *et al.* (2009) used chloroplast CAPS markers that differentiate between the chloroplasts of *B. napus* and *B. rapa* to survey wild and weedy populations of *B. rapa* for the capture of *B. napus* chloroplasts.

Intraspecific variation: Intraspecific diversity in plant cpDNA has been believed to be the result of mutations, genetic drift, selection by man for particular cpDNA-encoded characters (herbicide resistance), or the introgression of a foreign cpDNA (chloroplast capture) from related species during the early domestication of the taxon and even in wild taxa (Harris and Ingram 1991). Such introgression has been reported to be common for many plant species (e.g. in *Iris* species; Arnold *et al.* 1992). Systematics and phylogenetic implications of intraspecific variation that may be detected in cpDNA of plant species are reviewed by Soltis *et al.* (1992).

Prentice *et al.* (2003) implied chloroplast universal primers to study variation in cultivated and wild *Prunus avium*. Other examples for the use of universal primers to assess intraspecific variation are those carried out on *Argania spinosa* (*Sapotaceae*) of Morocco (El Mousadik and Petit 1996), *Primula cuneifolia* (*Primulaceae*) (Fujii *et al.* 1999), *Cannabis sativa* (Kohjyouma *et al.* 2000), the rare Spanish endemic, *Silene hifacensis* (Prentice *et al.* 2003), *Sagittaria latifolia* (Dorken and Barrett 2004), and *Oryza sativa* (Ou *et al.* 2009).

Identification of plants: There are several features of the chloroplast genome that make it well suited for species diagnosis. First, the low evolutionary rate of cpDNA below the species level, as was revealed by several studies (e.g. Jordan *et al.* 1996, Szalanski *et al.* 2001) means that sequence variation discovered between species is unlikely to be confounded by intraspecific variation at the same site. Second, the level of genetic variation in the chloroplast genome has been reported to be sufficient (Nissen *et al.* 1995, Lakshmi *et al.* 2002) to allow discrimination of closely related plant species. Such variation was even found to be numerous in some groups such as the *Senecio* complex (Bain and Jansen 1996), and species belonging to the *Solanaceae* (Bryan *et al.* 1999).

Linacre and Thorpe (1998) used DNA-specific primers for the nucleotide sequences between the *trnL* and *trnF* genes in the chloroplast genome for the unambiguous identification of *Cannabis sativa*. Similarly, Ohyama *et al.* (1999) used sequences of *trnL-trnF* and *trnT-trnL* to distinguish *Quercus gilva* from five other species in the genus *Quercus*. Brunner *et al.* (2001) was also able to identify fine roots of tree species of the Alps. Other examples are provided by Parani *et al.* (2001), who distinguished seven small millet species using PCR-RFLP on amplicons generated using universal primers that amplify the chloroplast *trnS-psbC* region (Demesure *et al.* 1995). Further, Fineschi *et al.* (2002), used *trnL* (and ITS2) sequences to differentiate between *Zelkova abelicea* and *Z. sicula* (*Ulmaceae*), and Haider and Nabulsi (2008), were able to identify *Aegilops* species based on restriction and sequencing of chloroplast loci, which they amplified using universal primers. More recently, universal primers have been proved to be of a great value for the DNA-barcoding (analysis of a short DNA sequence from a uniform target gene to enable species identification) of plant species (Kress and Erickson 2007, Ford *et al.* 2009).

Examples on other applications of universal primers in molecular studies on plant species are 1) analysis of phyto-geographic structure in plant species such as those of *Lolium* (Balfourier *et al.* 2000) and *Dryas integrifolia* (Tremblay and Schoen 1999), 2) investigation of the postglacial colonization history the plant species as the oak species *Quercus robur* and *Q. petraea* (Ferris *et al.* 1998), and 3) to determine the nucleotide sequence of a chloroplast region in a species of interest when that sequence is not available on public databases. For instance, Linacre and Thorpe (1998) used *trnL-F* universal primers (Taberlet *et al.* 1991) to recover the nucleotide sequence of this region in *Cannabis sativa*.

Conclusions

Universal primers have been developed to target the different three genomes (nuclear, mitochondrial and the chloroplastic) in plants. The most useful among these genomes, however, for the design of such primers has been the chloroplast genome. In certain occasions, and for some parts of the chloroplast genome, however, the design of chloroplast-specific universal primers is troublesome. This difficulty is experienced for chloroplast regions that have a higher rate of sequence evolution (intergenic spacers and introns), and when genes order rearrangements such as inversions, insertions, deletions, inverted repeats expansion or contraction or loss, and transpositions occur especially above the genus level in some plant groups.

Variation that can be detected among plant species using cpDNA-specific universal primers can be useful for many applications in plant science. Among these, systematics and evolutionary relationships studies at the different taxonomic levels (e.g. interfamilial, intergeneric, interspecific, intraspecific and interpopulational level)

have been by far the most prevalent. Examples of other studies that made use of such primers have been reported in this review.

The availability of such primers facilitates especially those studies that are usually carried out on a large set of different plant species (such as identification or DNA-barcoding studies). The availability of universal primers for molecular systematic studies on plants can save time, effort and cost that would otherwise be required to identify and sequence a suitable locus across all target plants.

Since the same genes in cpDNA may have different levels of variation among different groups of plant species and different levels of variation can be observed among the different genes of cpDNA within the same group, designing more universal primers that target both coding and non-coding regions would ease the choice of the chloroplast region to be targeted in molecular studies on a particular group, for a particular application and a particular taxonomic level.

References

- Angioi, S.A., Desiderio, F., Rau, D., Bitocchi, E., Attene, G., Rapa, R.: Development and use of chloroplast microsatellites in *Phaseolus* spp. and other legumes. - *Plant Biol.* **11**: 598-612, 2009.
- Aoki, K., Matsumura, T., Hattori, T., Murakami, N.: Chloroplast DNA phylogeography of *Photinia glabra* (*Rosaceae*) in Japan. - *Amer. J. Bot.* **93**: 1852-1858, 2006.
- Appels, R., Honeycutt, R.L.: rDNA: evolution over a billion years. - In: Dutta, S.K. (ed.): *DNA Systematics*. Vol. II, Plants. CRC Press, Boca Raton 1986.
- Arnold, M.L., Robinson, J.J., Buckner, C.M., Bennet, B.D.: Pollen dispersal and interspecific gene flow in Louisiana irises. - *Heredity* **68**: 399-404, 1992.
- Bain, J.F., Jansen, R.K.: Numerous chloroplast DNA polymorphisms are shared among different populations and species in the aureoid *Senecio* (Packera) complex. - *Can. J. Bot.* **74**: 1719-1728, 1996.
- Baldwin, B.G., Markos, S.: Phylogenetic utility of the external transcribed spacer (ETS) of 18S-26S rDNA: congruence of ETS and ITS trees of *Calycadenia* (*Compositae*). - *Mol. Phylogenet. Evol.* **10**: 449-463, 1998.
- Baldwin, B.G., Sanderson, M.J., Porter, J.M., Wojciechowski, M.F., Campbell, C.S., Donoghue, M.J.: The ITS region of nuclear ribosomal DNA – a valuable source of evidence on angiosperm phylogeny. - *Ann. Missouri Bot. Garden* **82**: 247-277, 1995.
- Balfourier, F., Imbert, C., Charmet, G.: Evidence for phylogeographic structure in *Lolium* species related to the spread of agriculture in Europe. A cpDNA study. - *Theor. appl. Genet.* **101**: 131-138, 2000.
- Bartish, I.V., Rumpunen, K., Nybom, H.: Combined analyses of RAPDs, cpDNA and morphology demonstrate spontaneous hybridisation in the plant genus *Chaenomeles*. - *Heredity* **85**: 383-392, 2000.
- Basha, S.D., Sujatha, M.: Genetic analysis of *Jatropha* species and interspecific hybrids of *Jatropha curcas* using nuclear and organelle specific markers. - *Euphytica* **168**: 197-214, 2009.
- Bastia, T., Carotenuto, N., Basile, B., Zoina, A., Cardi, T.: Induction of novel organelle DNA variation and transfer of resistance to frost and *Verticillium* wilt in *Solanum tuberosum* through somatic hybridization with *1EBN S. commersonii*. - *Euphytica* **116**: 1-10, 2000.
- Bastia, T., Scotti, N., Cardi, T.: Organelle DNA analysis of *Solanum* and *Brassica* somatic hybrids by PCR with 'universal primers'. - *Theor. appl. Genet.* **102**: 1265-1272, 2001.
- Bever, R., Brinkae, L., Jones, E.: Identification of plant trace evidence using molecular techniques. (<http://www.promega.com/geneticidproc/ussymp11proc/abstracts/bever.pdf>).
- Bever, R., Cimino, M.: *Forensic Molecular Botany: Identification of Plants from Trace Evidence*. - The Bode Technology Group, Springfield, 2001.
- Bobola, M.S., Eckert, R.T., Klein, A.S., Stapelfeldt, K., Smith, D.E., Guenette, D.: Using nuclear and organelle DNA markers to discriminate among *Picea rubens*, *Picea mariana*, and their hybrids. - *Can. J. Forest Res.* **26**: 433-443, 1996.
- Bouhadida, M.: Chloroplast DNA diversity in *Prunus* and its implication on genetic relationships. - *J. amer. Soc. hort. Sci.* **132**: 670-679, 2007.
- Brouat, C., Gielly, L., McKey, D.: Phylogenetic relationships in the genus *Leonardoxa* (*Leguminosae: Caesalpinioideae*) inferred from chloroplast *trnL* intron and *trnL-trnF* intergenic spacer sequences. - *Amer. J. Bot.* **88**: 143-149, 2001.
- Brunner, I., Brodbeck, S., Buchler, U., Sperisen, C.:

- Molecular identification of fine roots of trees from the Alps: reliable and fast DNA extraction and PCR-RFLP analyses of plastid DNA. - *Mol. Ecol.* **10**: 2079-2087, 2001.
- Bryan, G.J., McNicoll, J., Ramsay, G., Meyer, R.C., De Jong, W.S.: Polymorphic simple sequence repeat markers in chloroplast genomes of *Solanaceous* plants. - *Theor. appl. Genet.* **99**: 859-867, 1999.
- Buso, G.S.C., Rangel, P.H.N., Ferreira, M.E.: Analysis of random and specific sequences of nuclear and cytoplasmic DNA in diploid and tetraploid American wild rice species (*Oryza* spp.). - *Genome* **44**: 476-494, 2001.
- Cato, S.A., Richardson, T.E.: Inter- and intraspecific polymorphism at chloroplast SSR loci and the inheritance of plastids in *Pinus radiata* D Don. - *Theor. appl. Genet.* **93**: 587-592, 1996.
- Chat, J., Jauregui, B., Petit, R.J., Nadot, S.: Reticulate evolution in kiwifruit (*Actinidia*, *Actinidiaceae*) identified by comparing their maternal and paternal phylogenies. - *Amer. J. Bot.* **91**: 736-747, 2004.
- Cheng, Y.J., Guo, W.W., Deng, X.X.: Molecular characterization of cytoplasmic and nuclear genomes in phenotypically abnormal Valencia orange (*Citrus sinensis*) plus Meiwa kumquat (*Fortunella crassifolia*) intergeneric somatic hybrids. - *Plant Cell Rep.* **21**: 445-451, 2003.
- Cheng, Y.J., Meng, H.J., Guo, W.W., Deng, X.X.: Universal chloroplast primer pairs for simple sequence repeat analysis in diverse genera of fruit crops. - *J. Hort. Sci. Biotechnol.* **81**: 132-138, 2006.
- Chiang, T.Y., Schaal, B.A., Peng, C.I.: Universal primers for amplification and sequencing a noncoding spacer between the *atpB* and *rbcL* genes of chloroplast DNA. - *Bot. Bull. Acad. sin.* **39**: 245-250, 1998.
- Chung, S., Staub, J.E.: The development and evaluation of consensus chloroplast primer pairs that possess highly variable sequence regions in a diverse array of plant taxa. - *Theor. appl. Genet.* **107**: 757-767, 2003.
- Clegg, M.T.: Chloroplast gene-sequences and the study of plant evolution. - *PNAS* **90**: 363-367, 1993.
- Collins, D., Mill, R.R., Moller, M.: Species separation of *Taxus baccata*, *T. canadensis*, and *T. cuspidata* (*Taxaceae*) and origins of their reputed hybrids inferred from RAPD and cpDNA data. - *Amer. J. Bot.* **90**: 175-182, 2003.
- Cox, A.V., Bennett, M.D., Dyer, T.A.: Use of the polymerase chain reaction to detect spacer size heterogeneity in plant 5S-ribosomal-RNA gene clusters and to locate such clusters in wheat (*Triticum aestivum* L.). - *Theor. appl. Genet.* **83**: 684-690, 1992.
- Csencsics, D., Angelonea, S., Panigab, M., Rotachb, P., Rudowb, A., Sabiteb, E., Schwabb, P., Wohlhauserb, P., Holderegger, R.: A large scale survey of *Populus nigra* presence and genetic introgression from non-native poplars in Switzerland based on molecular identification. - *J. Natur. Conserv.* **17**: 142-149, 2009.
- Cubas, P., Pardo, C., Tahiri, H.: Genetic variation and relationships among *Ulex* (Fabaceae) species in southern Spain and northern Morocco assessed by chloroplast microsatellite (cpSSR) markers. - *Amer. J. Bot.* **92**: 2031-2043, 2005.
- Curtis, S.E., Clegg, M.T.: Molecular evolution of chloroplast DNA sequences. - *Mol. Biol. Evol.* **1**: 291-301, 1984.
- Dane, F., Lang, P., Bakhtiyarova, R.: Comparative analysis of chloroplast DNA variability in wild and cultivated *Citrullus* species. - *Theor. appl. Genet.* **108**: 958-966, 2004.
- De Lange, P.J., Heenan, P.B., Keeling, D.J., Murray, B.G., Smitsen, R., Sykes, W.R.: Biosystematics and conservation: a case study with two enigmatic and uncommon species of *Crassula* from New Zealand. - *Ann. Bot.* **101**: 881-99, 2008.
- Demesure, B., Comps, B., Petit, R.J.: Chloroplast DNA phylogeography of the common beech (*Fagus sylvatica* L.) in Europe. - *Evolution* **50**: 2515-2520, 1996.
- Demesure, B., Sodji, N., Petit, R.J.: A set of universal primers for amplification of polymorphic non-coding regions of mitochondrial and chloroplast DNA in plants. - *Mol. Ecol.* **4**: 129-131, 1995.
- Dorken, M.E., Barrett, S.C.H.: Chloroplast haplotype variation among monoecious and dioecious populations of *Sagittaria latifolia* (*Alismataceae*) in eastern North America. - *Mol. Ecol.* **13**: 2699-2707, 2004.
- Dumolin-Lapègue, S., Demesure, B., Le Corre V., Fineschi, S., Petit, R.J.: Phylogeographic structure of white oaks throughout the European continent. - *Genetics* **146**: 1475-1487, 1997b.
- Dumolin-Lapègue, S., Pemonge, M.H., Petit, R.J.: An enlarged set of consensus primers for the study of organelle DNA in plants. - *Mol. Ecol.* **6**: 393-398, 1997a.
- Eber, F., Tanguy, X., Chevre, A.M., Baranger, A., Vallee, P., Renard, M.: Spontaneous hybridisation between a male-sterile oilseed rape and two weeds. - *Theor. appl. Genet.* **88**: 362-368, 1994.
- El Mousadik, A., Petit, R.J.: Chloroplast DNA phylogeography of the argan tree of Morocco. - *Mol. Ecol.* **5**: 547-555, 1996.
- Ellstrand, N.C., Prentice, H.C., Hancock, J.F.: Gene flow and introgression from domesticated plants into their wild relatives. - *Ann. Rev. Ecol. Syst.* **30**: 539-563, 1999.
- Fant, J.B., Kamau, E.A., Preston, C.D.: Chloroplast evidence for the multiple origins of the hybrid *Potamogeton × sudermanicus* Hagstr. - *Aquat. Bot.* **75**: 351-356, 2003.
- Feliner, G.N., Aguilar, J.F., Rossello, J.A.: Reticulation or divergence: the origin of a rare serpentine endemic assessed with chloroplast, nuclear and RAPD markers. - *Plant Syst. Evol.* **231**: 19-38, 2002.
- Ferri, G., Alù, M., Corradini, B., Beduschi, G.: Forensic botany: species identification of botanical trace evidence using a multigene barcoding approach. - *Int. J. Legal Med.* **123**: 395-401, 2009.
- Ferris, C., King, R.A., Vainola, R., Hewitt, G.M.: Chloroplast DNA recognizes three refugial sources of European oaks and suggests independent eastern and western immigrations to Finland. - *Heredity* **80**: 584-593, 1998.
- Fineschi, S., Anzidei, M., Cafasso, D., Cozzolino, S., Garfi, G., Pastorelli, R., Salvini, D., Turchini, D., Vendramin, G.G.: Molecular markers reveal a strong genetic differentiation between two European relic tree species: *Zelkova abelicea* (Lam.) Boissier and *Z. sicula* Di Pasquale, Garfi and Quezel (*Ulmaceae*). - *Conserv. Genet.* **3**: 145-153, 2002.
- Fofana, B., Baudoin, J.P., Vekemans, X., Debouck, D.G., du Jardin, P.: Molecular evidence for an Andean origin and a secondary gene pool for the lima bean (*Phaseolus lunatus*

- L.) using chloroplast DNA. - *Theor. appl. Genet.* **98**: 202-212, 1999.
- Fofana, B., Harvengt, L., Baudoin, J.P., DuJardin, P.: New primers for the polymerase chain amplification of cpDNA intergenic spacers in *Phaseolus* phylogeny. - *Belg. J. Bot.* **129**: 118-122, 1997.
- Ford, C.S., Ayres, K.L., Haider, N., Toomey, N., Stohl, J.V.A., Kelly, L., Wilstöm, N., Hollingsworth, P., Duff, R.J., Hoot, S.B., Cowan, R.S., Chase, M.W., Wilkinson, M.J.: Selection of candidate DNA barcoding regions for use on land plants. - *Bot. J. Linn. Soc.* **159**: 1-11, 2009.
- Friesen, N., Fritsch, R.M., Pollner, S., Blattner, F.R.: Molecular and morphological evidence for an origin of the aberrant genus *Milula* within Himalayan species of *Allium* (*Alliaceae*). - *Mol. Phylogenet. Evol.* **17**: 209-218, 2000.
- Friesen, N., Fritsch, R.M., Pollner, S., Blattner, F.R.: RAPDs and noncoding chloroplast DNA reveal a single origin of the cultivated *Allium fistulosum* from *A. altaicum* (*Alliaceae*). - *Amer. J. Bot.* **86**: 554-562, 1999.
- Fu, C.H., Chen, C.L., Guo, W.W., Deng, X.X.: GISH, AFLP and PCR-RFLP analysis of an intergeneric somatic hybrid combining goutou sour orange and *Poncirus trifoliata*. - *Plant Cell Rep.* **23**: 391-396, 2004.
- Fujii, N., Ueda, K., Watano, Y., Shimizu, T.: Further analysis of intraspecific sequence variation of chloroplast DNA in *Primula cuneifolia* Ledeb. (*Primulaceae*): implications for biogeography of the Japanese alpine flora. - *J. Plant Res.* **112**: 87-95, 1999.
- Gillespie, L.J., Boles, R.: Phylogenetic relationships and infraspecific variation in Canadian arctic *Poa* based on chloroplast DNA restriction site data. - *Can. J. Bot.* **79**: 679-701, 2001.
- Gillespie, L.J., Soreng, R.J.: A phylogenetic analysis of the bluegrass genus *Poa* based on cpDNA restriction site data. - *Syst. Bot.* **30**: 84-105, 2005.
- Goremykin, V., Bobrova, V., Pahnke, J., Troitsky, A., Antonov, A., Martin, W.: Noncoding sequences from the slowly evolving chloroplast inverted repeat in addition to *rbcL* data do not support Gnetales affinities of angiosperms. - *Mol. Biol. Evol.* **13**: 383-396, 1996.
- Graham, S.W., Olmstead, R.G.: Utility of 17 chloroplast genes for inferring the phylogeny of the basal angiosperms. - *Amer. J. Bot.* **87**: 1712-1730, 2000.
- Griffin, S.R., Barrett, S.C.H.: Post-glacial history of *Trillium grandiflorum* (*Melanthiaceae*) in Eastern North America: Inferences from phylogeography. - *Amer. J. Bot.* **91**: 465-473, 2004.
- Grivet, D., Heinze, B., Vendramin, G.G., Petit, R.J.: Genome walking with consensus primers: application to the large single copy region of chloroplast DNA. - *Mol. Ecol. Notes* **1**: 345-349, 2001.
- Haider, N.: Development and use of universal primers in plants. - Ph.D. Thesis, University of Reading, Reading 2003.
- Haider, N., Allainguillaume, J., Wilkinson, M.J.: The inefficiency of chloroplast transformation in preventing transgene flow: a case study of chloroplast capture. - *Curr. Genet.* **55**: 139-50, 2009.
- Haider, N., Nabulsi, I.: Identification of *Aegilops* L. species and *Triticum aestivum* L. based on chloroplast DNA. - *Genet. Resour. Crop Evol.* **55**: 537-549, 2008.
- Haider, N., Wilkinson, M.J.: A set of plastid DNA-specific universal primers for flowering plants. - Submitted to *Russ. J. Genet.*
- Hamilton, M.B.: Four primer pairs for the amplification of chloroplast intergenic regions with intraspecific variation. - *Mol. Ecol.* **8**: 521-523, 1999.
- Harris, S.A., Ingram, R.: Chloroplast DNA and biosystematics - the effects of intraspecific diversity and plastid transmission. - *Taxon* **40**: 393-412, 1991.
- Hollingsworth, M.L., Bailey, J.P., Hollingsworth, P.M., Ferris, C.: Chloroplast DNA variation and hybridization between invasive populations of Japanese knotweed and giant knotweed (*Fallopia, Polygonaceae*). - *Bot. J. Linn. Soc.* **129**: 139-154, 1999.
- Honjo, M., Ueno, S., Tsumura, Y., Washitani, I., Ohsawa, R.: Phylogeographic study based on intraspecific sequence variation of chloroplast DNA for the conservation of genetic diversity in the Japanese endangered species *Primula sieboldii*. - *Biol. Conserv.* **120**: 211-220, 2004.
- Horres, R., Zizka, G., Kahl, G., Weising, K.: Molecular phylogenetics of *Bromeliaceae*: evidence from *trnL* (UAA) intron sequences of the chloroplast genome. - *Plant Biol.* **2**: 306-315, 2000.
- Hu, D., Zhang, Q., Luo, Z.: Phylogenetic analysis in some *Diospyros* spp. (*Ebenaceae*) and Japanese persimmon using chloroplast DNA PCR-RFLP markers. - *Scientia Hort.* **117**: 32-38, 2008.
- Jordan, W.C., Courtney, M.W., Neigel, J.E.: Low levels of intraspecific genetic variation at a rapidly evolving chloroplast DNA locus in North American duckweeds (*Lemnaceae*). - *Amer. J. Bot.* **83**: 430-439, 1996.
- King, R.A., Ferris, C.: Chloroplast DNA phylogeography of *Alnus glutinosa* (L.) Gaertn. - *Mol. Ecol.* **7**: 1151-1161, 1998.
- Kohjyouma, M., Lee, I.J., Iida, O., Kurihara, K., Yamada, K., Makino, Y., Sekita, S., Satake, M.: Intraspecific variation in *Cannabis sativa* L. based on intergenic spacer region of chloroplast DNA. - *Biol. Pharm. Bull.* **23**: 727-730, 2000.
- Kress, W.J., Erickson, D.L.: A two-locus global DNA barcode for land plants: the coding *rbcL* gene complements the non-coding *trnH-psbA* spacer region. - *PLoS ONE* **2**: e508, 2007.
- Lakshmi, M., Parani, M., Parida, A.: Molecular phylogeny of mangroves IX. - Molecular marker assisted intra-specific variation and species relationships in the Indian mangrove tribe *Rhizophoraceae*. - *Aquat. Bot.* **74**: 201-217, 2002.
- Lavigne, C., Klein, E.K., Couvet, D.: Using seed purity data to estimate an average pollen mediated gene flow from crops to wild relatives. - *Theor. appl. Genet.* **104**: 139-145, 2002.
- Lee, C., Wen, J.: Phylogeny of *Panax* using chloroplast *trnC-trnD* intergenic region and the utility of *trnC-trnD* in interspecific studies of plants. - *Mol. Phylogenet. Evol.* **31**: 894-903, 2004.
- Lihová, J., Kuera, J., Pern, M., Marhold, K.: Hybridization between two polyploid *Cardamine* (*Brassicaceae*) species in North-Western Spain: discordance between morphological and genetic variation patterns. - *Ann. Bot.* **99**: 1083-1096, 2007.
- Linacre, A., Thorpe, J.: Detection and identification of cannabis by DNA. - *Forensic Sci. Int.* **91**: 71-76, 1998.
- Linder, C.R., Moore, L.A., Jackson, R.B.: A universal

- molecular method for identifying underground plant parts to species. - *Mol. Ecol.* **9**: 1549-1559, 2000.
- Lotfy, S., Luro, F., Carreel, F., Froelicher, Y., Rist, D., Ollitrault, P.: Application of cleaved amplified polymorphic sequence method for analysis of cytoplasmic genome among *Aurantioideae* intergeneric somatic hybrids. - *J. amer. Soc. hort. Sci.* **128**: 225-230, 2003.
- Matsushima, R., Hu, Y., Toyoda, K., Sodmergen, Sakamoto, W.: The model plant *Medicago truncatula* exhibits biparental plastid inheritance. - *Plant Cell Physiol.* **49**: 81-91, 2008.
- Matyas, G., Sperisen, C.: Chloroplast DNA polymorphisms provide evidence for postglacial re-colonisation of oaks (*Quercus spp.*) across the Swiss Alps. - *Theor. appl. Genet.* **102**: 12-20, 2001.
- McIntosh, S.R., Pacey-Miller, T., Henery, R.J.: A universal protocol for identification of cereals. - *J. Cereal Sci.* **41**: 37-46, 2005.
- Meerow, A.W., Lehmiller, D.J., Clayton, J.R.: Phylogeny and biogeography of *Crinum* L. (*Amaryllidaceae*) inferred from nuclear and limited plastid non-coding DNA sequences. - *Bot. J. Linn. Soc.* **141**: 349-363, 2003.
- Meudt, H.M., Bayly, M.J.: Phylogeographic patterns in the Australasian genus *Chionohebe* (Veronica s.l., *Plantaginaceae*) based on AFLP and chloroplast DNA sequences. - *Mol. Phylogenet. Evol.* **47**: 319-38, 2008.
- Moore, L.A., Field, C.B.: A technique for identifying the roots of different species in mixed samples using nuclear ribosomal DNA. - *J. veg. Sci.* **16**: 131-134, 2005.
- Natali, A., Manen, J.F., Ehrendorfer, F.: Phylogeny of the *Rubiaceae-Rubioideae*, in particular the tribe *Rubieae*: evidence from a noncoding chloroplast DNA sequence. - *Ann. Missouri Bot. Garden* **82**: 428-439, 1995.
- Nishikawa, T., Salomon, B., Komatsuda, T., Von Bothmer, R., Kadowaki, K.: Molecular phylogeny of the genus *Hordeum* using three chloroplast DNA sequences. - *Genome* **45**: 1157-1166, 2002.
- Nishizawa, T., Watano, Y.: Primer pairs suitable for PCR-SSCP analysis of chloroplast DNA in angiosperms. - *J. Phytoeogr. Taxon.* **48**: 63-66, 2000.
- Nissen, S.J., Masters, R.A., Lee, D.J., Rowe, M.L.: DNA-based marker systems to determine genetic diversity of weedy species and their application to biocontrol. - *Weed Sci.* **43**: 504-513, 1995.
- Oddou-Muratorio, S., Petit, R.J., Le Guerroue, B., Guesnet, D., Demesure, B.: Pollen- versus seed-mediated gene flow in a scattered forest tree species. - *Evolution* **55**: 1123-1135, 2001.
- Ofori, D.A., Swaine, M.D., Leifert, C., Cobbinah, J.R., Price, A.H.: Population genetic structure of *Milicia* species characterised by using RAPD and nucleotide sequencing. - *Genet. Resour. Crop Evol.* **48**: 637-647, 2001.
- Ohyama, M., Baba, K., Itoh, T.: Possibility of grouping of *Cyclobalanopsis* species (*Fagaceae*) grown in Japan based on an analysis of several regions of chloroplast DNA. - *J. Wood Sci.* **45**: 498-501, 1999.
- Oliver, C., Hollingsworth, P.M., Gornall, R.J.: Chloroplast DNA phylogeography of the arctic-montane species *Saxifraga hirculus* (*Saxifragaceae*). - *Heredity* **96**: 222-231, 2006.
- Olivares-Fuster, Ó., Hernández-Garrido, M., Guerri, J., Navarro, L.: Plant somatic hybrid cytoplasmic DNA characterization by single-strand conformation polymorphism. - *Tree Physiol.* **27**: 785-792, 2007.
- Olmstead, R.G., Palmer, J.D.: Chloroplast DNA systematics - a review of methods and data-analysis. - *Amer. J. Bot.* **81**: 1205-1224, 1994.
- Ou, L.J., Huang, G.W., Li, W.J., Kang, G.P., Chen, J.L., Luan, S., Chen, L.B.: Chloroplast DNA polymorphism in different types of cytoplasmic male sterile rice. - *Biol. Plant.* **53**: 593-596, 2009.
- Palmer, J.D.: Chloroplast DNA evolution and biosystematic uses of chloroplast DNA variation. - *Amer. Natur.* **130** (Suppl.): S6-S29, 1987.
- Pan, Y., Shi, S., Gong, X., Kuroda, C.: A natural hybrid between *Ligularia paradoxa* and *L. duciformis* (*Asteraceae, Senecioneae*) from Yunnan, China. - *Ann. Missouri Bot. Garden* **95**: 487-494, 2008.
- Panda, S., Martin, J.P., Aguinalalde, I.: Chloroplast DNA study in sweet cherry cultivars (*Prunus avium* L.) using PCR-RFLP method. - *Genet. Resour. Crop Evol.* **50**: 489-495, 2003.
- Parani, M., Rajesh, K., Lakshmi, M., Parducci, L., Szmidi, A.E., Parida, A.: Species identification in seven small millet species using polymerase chain reaction-restriction fragment length polymorphism of *trnS-psbC* gene region. - *Genome* **44**: 495-499, 2001.
- Parducci, L., Szmidi, A.E.: PCR-RFLP analysis of cpDNA in the genus *Abies*. - *Theor. appl. Genet.* **98**: 802-808, 1999.
- Petit, R.J., Demesure, B., Pineau, E.: Chloroplast DNA differentiation at a local and continental scale in European oak species: the importance of historical factors. - In: Kremer, A., Muhs, H.J. (ed.): *Inter- and Intraspecific Variation in European Oaks: Evolutionary Implications and Practical Consequences*. Pp. 145-164. Office for Official Publications of the European Communities, Luxembourg 1996.
- Petit, R.J., Pineau, E., Demesure, B., Bacilieri, R., Ducousso, A., Kremer, A.: Chloroplast DNA footprints of postglacial recolonization by oaks. - *PNAS* **94**: 9996-10001, 1997.
- Pilepic, K.H.: Analysis of chloroplast DNA variability among some *Hypericum* species in Croatia. - *Period. Biol.* **104**: 457-462, 2002.
- Potgieter, K., Albert, V.A.: Phylogenetic relationships within *Apocynaceae* s.l. based on *trnL* intron and *trnL-F* spacer sequences and propagule characters. - *Ann. Missouri Bot. Garden* **88**: 523-549, 2001.
- Prado, M.D., De Prado, R., Franco, A.R.: Design and optimization of degenerated universal primers for the cloning of the plant acetolactate synthase conserved domains. - *Weed Sci.* **52**: 487-491, 2004.
- Prentice, H.C., Malm, J.U., Mateu-Andres, I., Segarra-Moragues, J.G.: Allozyme and chloroplast DNA variation in island and mainland populations of the rare Spanish endemic, *Silene hifacensis* (*Caryophyllaceae*). - *Conserv. Genet.* **4**: 543-555, 2003.
- Provan, J., Biss, P.M., McMeel, D., Mathews, S.: Universal primers for the amplification of chloroplast microsatellites in grasses (*Poaceae*). - *Mol. Ecol. Notes* **4**: 262-264, 2004a.
- Provan, J., Corbett, G., McNicol, J.W., Powell, W.: Chloroplast DNA variability in wild and cultivated rice (*Oryza spp.*) revealed by polymorphic chloroplast simple sequence repeats. - *Genome* **40**: 104-110, 1997.

- Provan, J., Murphy, S., Maggs, C.A.: Universal plastid primers for *Chlorophyta* and *Rhodophyta*. - *Eur. J. Phycol.* **39**: 43-50, 2004b.
- Provan, J., Powell, W., Hollingsworth, P.M.: Chloroplast microsatellites: new tools for studies in plant ecology and evolution. - *Trends Ecol. Evol.* **16**: 142-147, 2001.
- Rebound, X., Zeyl, C.: Organelle inheritance in plants. - *Heredity* **72**: 132-140, 1994.
- Samigullin, T.H., Valiejo-Roman, K.M., Troitsky, A.V., Bobrova, V.K., Filin, V.R., Martin, W., Antonov, A.S.: Sequences of rDNA internal transcribed spacers from the chloroplast DNA of 26 bryophytes: properties and phylogenetic utility. - *FEBS Letters* **422**: 47-51, 1998.
- Sass, C., Little, D.P., Stevenson, D.W., Specht, C.D.: DNA barcoding in the *cycadales*: testing the potential of proposed barcoding markers for species identification of cycads. - *PLoS ONE* **2**: e1154, 2007.
- Soltis, D.E., Soltis, P.S., Milligan, B.G.: Intraspecific chloroplast DNA variation: systematics and phylogenetic implications. - In: Soltis, P.S., Soltis, D.E., Doyle, J.J. (ed.): *Molecular Systematics of Plants*. Pp. 117-150. Chapman and Hall, New York 1992.
- Sperisen, C., Gugerli, F., Büchler, U., Mátyás, G.: Comparison of two rapid DNA extraction protocols for gymnosperms for application in population genetic and phylogenetic studies. - *Forest Genet.* **7**: 133-136, 2000.
- Sun, G.L.: Interspecific polymorphism at non-coding regions of chloroplast, mitochondrial DNA and rRNA IGS region in *Elymus* species. - *Hereditas* **137**: 119-124, 2002.
- Sun, Y., Skinner, D.Z., Liang, G.H., Hulbert, S.H.: Phylogenetic analysis of sorghum and related taxa using internal transcribed spacers of nuclear ribosomal DNA. - *Theor. appl. Genet.* **89**: 26-32, 1994.
- Szalanski, A.L., Steinauer, G., Bischof, R., Petersen, J.: Origin and conservation genetics of the threatened Ute ladies'-tresses, *Spiranthes diluvialis* (*Orchidaceae*). - *Amer. J. Bot.* **88**: 177-180, 2001.
- Taberlet, P., Gielly, L., Pautou, G., Bouvet, J.: Universal primers for amplification of three non-coding regions of chloroplast DNA. - *Plant mol. Biol.* **17**: 1105-1109, 1991.
- Tanksley, S.D., Pichersky, E.: Organisation and evolution of sequences in the plant nuclear genome. - In: Gottlieb, L.D., Jain, S.K. (ed.): *Plant Evolutionary Biology*. Pp. 55-83. Chapman and Hall, London - New York 1988.
- Taylor, C., Madsen, K., Borg, S., Moller, M.G., Boelt, B., Holm, P.B.: The development of sequence-tagged sites (STSs) in *Lolium perenne* L.: the application of primer sets derived from other genera. - *Theor. appl. Genet.* **103**: 648-658, 2001.
- Terry, R.G., Nowak, R.S., Tausch, R.J.: Genetic variation in chloroplast and nuclear ribosomal DNA in Utah juniper (*Juniperus osteosperma*, *Cupressaceae*): evidence for interspecific gene flow. - *Amer. J. Bot.* **87**: 250-258, 2000.
- Testolin, R., Cipriani, G.: Paternal inheritance of chloroplast DNA and maternal inheritance of mitochondrial DNA in the genus *Actinidia*. - *Theor. appl. Genet.* **94**: 897-903, 1997.
- Tremblay, N.O., Schoen, D.J.: Molecular phylogeography of *Dryas integrifolia*: glacial refugia and postglacial recolonization. - *Mol. Ecol.* **8**: 1187-1198, 1999.
- Tsumura, Y., Kawahara, T., Wickneswari, R., Yoshimura, K.: Molecular phylogeny of *Dipterocarpaceae* in Southeast Asia using RFLP of PCR-amplified chloroplast genes. - *Theor. appl. Genet.* **93**: 22-29, 1996.
- Vakhitov, V.A., Chemeris, A.V., Sabirzhanov, B.E., Akhunov, E.D., Kulikov, A.M., Nikonorov, Y.M., Gimalov, F.R., Bikbulatova, S.M., Baymiev, A.K.: The phylogeny of *Triticum* L. and *Aegilops* L. inferred from comparative analysis of nucleotide sequences in rDNA promoter regions. - *Russ. J. Genet.* **39**: 1-11, 2003.
- Van Droogenbroeck, B., Maertens, I., Haegeman, A., Kyndt, T., O'Brien, C., Drew, R.A., Gheysen, G.: Maternal inheritance of cytoplasmic organelles in intergeneric hybrids of *Carica papaya* L. and *Vasconcellea* spp. (*Caricaceae* Dumort., *Brassicaceae*). - *Euphytica* **143**: 161-168, 2005.
- Vendramin, G.G., Lelli, L., Rossi, P., Morgante, M.: A set of primers for the amplification of 20 chloroplast microsatellites in *Pinaceae*. - *Mol. Ecol.* **5**: 595-598, 1996.
- Wang, B.C., Smith, T.B.: Closing the seed dispersal loop. - *Trends Ecol. Evol.* **17**: 379-385, 2002.
- Watts, C.D., Fisher, A.E., Shrum, C.D., Newbold, W.Y.L., Hansen, S., Liu, C., Scot, A.K.: The D4 set: primers that target highly variable intron loops in plant chloroplast genomes. - *Mol. Ecol. Resour.* **8**: 1344-1347, 2008.
- Weber, K.L., Bolander, M.E., Sarkar, G.: Rapid acquisition of unknown DNA sequence adjacent to a known segment by multiplex restriction site PCR. - *Biotechniques* **25**: 415-419, 1998.
- Weising, K., Gardner, R.C.: A set of conserved PCR primers for the analysis of Simple Sequence Repeat polymorphisms in chloroplast genomes of dicotyledonous angiosperms. - *Genome* **42**: 9-19, 1999.
- Wesselink, M., Kuiper, I.: Species identification of botanical trace evidence using molecular markers. - *Forensic Sci. Int.* **1** (Genet. Suppl.): 630-632, 2008.
- White, T.J., Bruns, T., Lee, S., Taylor, J.: Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. - In: Innis, M.A., Gelfand, D.H., Sninsky, J.J., White, T.J. (ed.): *PCR Protocols: a Guide to Methods and Applications*. Pp. 315-322. Academic Press, San Diego 1990.
- Wilkinson, M., Linacre, A.M.T.: The detection and persistence of *Cannabis sativa* DNA on skin. - *Sci. Justice* **40**: 11-14, 2000.
- Yang, T.W., Yang, Y.A., Xiong, Z.G.: Paternal inheritance of chloroplast DNA in interspecific hybrids in the genus *Larrea* (*Zygophyllaceae*). - *Amer. J. Bot.* **87**: 1452-1458, 2000.
- Yang, Y.W., Tai, P.Y., Chen, Y., Li, W.H.: A study of the phylogeny of *Brassica rapa*, *B. nigra*, *Raphanus sativus*, and their related genera using noncoding regions of chloroplast DNA. - *Mol. Phylogenet. Evol.* **23**: 268-275, 2002.
- Zhang, C., Fan, X., Yu, H.Q., Zhang, L., Wang, X.L., Zhou, Y.H.: Different maternal genome donor to *Kengyilia* species inferred from chloroplast *trnL-F* sequences. - *Biol. Plant.* **53**: 759-763, 2009.
- Ziegenhagen, B., Fladung, M.: Variation in the *psbC* gene region of gymnosperms and angiosperms as detected by a single restriction site polymorphism. - *Theor. appl. Genet.* **94**: 1065-1071, 1997.
- Zuber, D., Widmer, A.: Genetic evidence for host specificity in the hemi-parasitic *Viscum album* L. (*Viscaceae*). - *Mol. Ecol.* **9**: 1069-1073, 2000.