

BRIEF COMMUNICATION

Hydrogen sulfide is involved in the regulation of ascorbate and glutathione metabolism by jasmonic acid in *Arabidopsis thaliana*

C. SHAN^{1,2,3,4}, T. WANG^{3,4}, Y. ZHOU^{1,3,4*}, and W. WANG²Postdoctoral Research Base, Henan Institute of Science and Technology, Xinxiang, 453003 P.R. China¹Postdoctoral Research Centre, Henan Agricultural University, Zhengzhou, 450002 P.R. China²School of Science and Technology, Henan Institute of Science and Technology, Xinxiang, 453003 P.R. China³Collaborative Innovation Center of Modern Biological Breeding, Henan Province, Xinxiang, 453003 P.R. China⁴

Abstract

This study investigated the role of hydrogen sulfide (H₂S) in the regulation of the ascorbate (AsA) and glutathione (GSH) metabolism by jasmonic acid (JA) in the leaves of *Arabidopsis thaliana* by using H₂S scavenger hypotaurine (HT) and H₂S synthetic mutant (SALK_041918, designated *Atl-cdes*). The results showed that JA significantly increased the H₂S content, the activities of L-cysteine desulphydrase (L-CDes), D-cysteine desulphydrase (D-CDes), ascorbate peroxidase (APX), glutathione reductase (GR), monodehydroascorbate reductase (MDHAR), dehydroascorbate reductase (DHAR), L-galactono-1,4-lactone dehydrogenase (GalLDH) and γ -glutamylcysteine synthetase (γ -ECS), the ratio of AsA to dehydroascorbate (DHA), and decreased the content of malondialdehyde (MDA) and H₂O₂ in the wild type of *A. thaliana*, compared to control. The above effects of JA except the increased activities of L-CDes and D-CDes were suppressed by addition of HT. However, JA and HT+JA had no significant effects on the ratio of reduced GSH to oxidized GSH (GSSG) in the wild type of *A. thaliana*. Application of HT to the control decreased H₂S content, AsA/DHA ratio, and activities of APX, GR, DHAR, MDHAR, γ -ECS, and GalLDH, but had no effects on MDA content, activities of L-CDes and D-CDes, and GSH/GSSG ratio. In the H₂S synthetic mutant, JA had no obvious effects on above mentioned parameters except the D-CDes activity compared with the control. Our results suggest that JA-induced H₂S, which is a signal that leads to the up-regulation of the AsA and GSH metabolism.

Additional key words: ascorbate peroxidase, cysteine desulphydrase, glutathione reductase, H₂S synthetic mutant, hypotaurine.

Ascorbate (AsA) and glutathione (GSH) have important roles in defence against oxidative damage induced by stresses. Ascorbate-glutathione cycle is the regeneration pathway of AsA and GSH through ascorbate peroxidase (APX), monodehydroascorbate reductase (MDHAR), dehydroascorbate reductase (DHAR), and glutathione reductase (GR) (Noctor and Foyer 1998). Through this cycle, AsA and GSH are regenerated and H₂O₂ is

scavenged, which, in turn, alleviates oxidative damage induced by reactive oxygen species (ROS). L-galactono-1,4-lactone dehydrogenase (GalLDH) and γ -glutamylcysteine synthetase (γ -ECS) are the key enzymes which are responsible for the biosynthesis of AsA and GSH, respectively. The regeneration and biosynthesis pathway of AsA and GSH have important roles in regulating their redox states, which, in turn, regulates the responses of

Submitted 30 September 2016, last revision 12 March 2017, accepted 15 March 2017.

Abbreviations: APX - ascorbate peroxidase; AsA - ascorbate; DHAR - dehydroascorbate reductase; γ -ECS - γ -glutamylcysteine synthetase; GalLDH - L-galactono-1,4-lactone dehydrogenase; GR - glutathione reductase; GSH - glutathione; HT - hypotaurine; JA - jasmonic acid; MDA - malondialdehyde; MDHAR - monodehydroascorbate reductase; WT - wild type.

Acknowledgements: This study was funded by the Startup Foundation for Postdoctors of Postdoctoral Research Centre, Henan Agricultural University and Postdoctoral Research Base, Henan Institute of Science and Technology, the Scientific Research Foundation for Postdoctors of Henan Province (2015104), Project of Supporting Young Backbone Teachers of Colleges and Universities in Henan province (2016GGJS-108) and the Innovation Fund of Science and Technology of Henan Institute of Science and Technology in 2015.

* Corresponding author; e-mail: shchjuan1978@aliyun.com

plants to stresses. Thus, it is very important to study the regulatory mechanism of the redox state of AsA and GSH.

Jasmonic acid (JA) is an important plant hormone that plays important roles in regulating stress responses as well as plant growth and development (Radhika *et al.* 2010, Ren and Dai 2012). Increasing evidence shows that JA can enhance the metabolism of AsA and GSH in plants. Our previous studies (Shan *et al.* 2011a, 2015) showed that signal molecules NO and mitogen-activated protein kinase kinase 1/2 (MEK1/2) participate in signal transduction of JA during regulation of the AsA and GSH metabolism.

Hydrogen sulfide is the third gaseous signalling molecule following nitric oxide (NO) and carbon monoxide (CO). In plants, H₂S can promote root organogenesis (Zhang *et al.* 2009a) and seed germination (Zhang *et al.* 2010a). Increasing evidence has proven that exogenous H₂S affects the antioxidative response under heavy metal, drought, and salt stresses (Zhang *et al.* 2008, 2009b, 2010b, Li *et al.* 2012, Wang *et al.* 2012). Our previous study showed that exogenous H₂S could regulate the metabolism of AsA and GSH in wheat leaves under osmotic stress (Shan *et al.* 2011b). It has been reported that H₂S is involved in the JA-induced stomatal closure (Hou *et al.* 2011). However, whether endogenous H₂S participates in the regulation of the AsA and GSH metabolism by JA remains unknown. To fulfil this gap, the current study investigated the content of H₂S, malondialdehyde (MDA), and H₂O₂, and the activities of H₂S synthetases L-cysteine desulphydrase (L-CDes) and D-cysteine desulphydrase (D-CDes), and the enzymes involved in the ascorbate and glutathione metabolism in the leaves of *Arabidopsis thaliana* treated with JA.

The seeds of *Arabidopsis thaliana* L. ecotype Col-0 (used as wild-type, WT) were supplied by Arabidopsis Biological Resource Center of USA. The mutant seeds of *A. thaliana* (*Atl-cdes*) based on the same ecotype were supplied by Prof. Xin Liu of Qingdao Agricultural University. *Atl-cdes* is the T-DNA insertion mutant of H₂S synthetase L-CDes (Liu *et al.* 2011). In this mutant, the synthetic pathway through L-CDes is interrupted. The *A. thaliana* seedlings were grown in half-strength Hoagland's solution for 4 weeks and then transferred to the half-strength Hoagland's solution containing 30 µM JA. A group of plants was pretreated with H₂S scavenger hypotaurine (HT; Peng *et al.* 2016) by soaking roots in half-strength Hoagland's solution containing 20 µM HT for 8 h and then exposed to 30 µM JA for 48 h. Control plants were treated with half-strength Hoagland's solution alone. Also the H₂S synthesis mutant plants were treated by JA for 48 h. Each treatment consisted of five replicates with 30 plants each. Hoagland's solutions were replaced by corresponding fresh solutions every day. The top fully expanded leaves of plants were harvested after 24 and 48 h of JA-treatment, frozen in liquid nitrogen, and stored at -80 °C until analyses. All the experiments

were conducted in a greenhouse under a temperature of 22 ± 2 °C, a relative humidity of 70 %, an irradiance of 120 µmol(photons) m⁻² s⁻¹, and a 16-h photoperiod.

Enzymes were extracted according to Shan and Liang (2010). Ascorbate peroxidase (APX, EC 1.11.1.11) activity was measured according to Nakano and Asada (1981). One unit of enzyme was defined as the amount of APX catalyzing the oxidation of 1 µmol of ascorbate per minute. Glutathione reductase (GR, EC 1.6.4.2) activity was monitored at 340 nm in a reaction mixture containing 100 mM Tris-HCl (pH 8.0), 0.5 mM ethylenediamine-tetraacetic acid, 0.5 mM MgCl₂, 10 mM oxidized glutathione (GSSG), 1 mM NADPH, and enzyme extract. The reaction was initiated by adding NADPH (Grace and Logan 1996). One unit of GR activity was defined as the reduction of 1 µmol of NADPH per minute. Monodehydroascorbate reductase (MDHAR, EC 1.6.5.4) activity was assayed at 340 nm (Miyake and Asada 1992). One unit of MDHAR activity was defined as the amount of enzyme that oxidized 1 µmol of NADH per minute. Dehydroascorbate reductase (DHAR, EC 1.8.5.1) activity was measured at 265 nm in assay solution containing 100 mM Hepes-KOH (pH 7.0), 20 mM GSH, 2 mM dehydroascorbate (DHA). The reaction was initiated by adding DHA (Dalton *et al.* 1986). One unit of DHAR activity was defined as the amount of enzyme that produced 1 µmol of ascorbate (AsA) per minute. L-galactono-1,4-lactone dehydrogenase (GalLDH, EC 1.3.2.3) was extracted and measured according to Shan and Liang (2010). Frozen samples were ground in a prechilled mortar with ice-cold 0.1 M potassium phosphate buffer (pH 7.4) containing 0.4 M sucrose. The extract was passed through 2 layers of miracloth and then centrifuged at 300 g for 10 min. The supernatant was centrifuged at 10 000 g for 20 min. The sediment was suspended in the above buffer containing 0.4 M sucrose. The assay mixture was composed of enzyme solution, 1.05 mg cm⁻³ cytochrom *c* dissolved in potassium phosphate buffer, and 56 mM L-galactono-1,4-lactone (L-GalL). Before assay, the mixture was incubated at 25 °C for 1 min. The increase in absorbance at 550 nm was followed immediately after the addition of L-Gal. One unit of GalLDH activity was defined as the amount of enzyme required to oxidize 1 nmol of L-Gal per minute. A γ-glutamylcysteine synthetase (γ-ECS, EC 6.3.2.2) was extracted and measured according to Rüeggseger and Brunold (1992). Frozen samples were ground in prechilled mortar with 0.1 M precooled Tris-HCl (pH 8.0). The extract was centrifuged at 20 000 g for 10 min. The supernatant was used for the assay of enzyme activity. The mixture of supernatant, 0.1 M Tris-HCl (pH 8.0), 0.25 mM glutamate, 10 mM ATP, 1 mM dithioerythritol, and 2 mM cysteine reacted at 25 °C for 1 h. Then, phosphorus agent, which consisted of 3 mM H₂SO₄, distilled water, 2.5 % (m/v) ammonium molybdate, and 10 % (m/v) AsA, was added and mixed. The mixture was incubated at 45 °C for 25 min and

cooled at room temperature. Then, the absorbance at 660 nm was measured. One unit of γ -ECS activity was defined as 1 μ mol of cysteine-generated PO_4^{3-} per minute determined by the molybdenum blue method. A molar coefficient of $5.6 \text{ mM}^{-1} \text{ cm}^{-1}$ was used for the calculation of enzyme activity. The activities of *L*-cysteine desulfhydrase (*L*-CDes, EC 4.4.1.1) and *D*-cysteine desulfhydrase (*D*-CDes, EC 4.1.99.4) were measured according to the method of Riemenschneider *et al.* (2005). One unit of *L*-CDes and *D*-CDes activities was defined as the formation of 1 nmol of H_2S per minute. The specific activities of all above enzymes were expressed as units per mg of protein. Protein content was determined by the method of Bradford (1976).

The content of AsA and DHA was measured according to Hodges *et al.* (1996) based on the reduction of ferric to ferrous ion with AsA in acidic solution. The content of GSSG and GSH was measured according to Griffith (1980) based on the enzymatic recycling method and a thiol-masking reagent, 2-vinylpyridine. H_2S was determined by formation of methylene blue from dimethyl-*p*-phenylenediamine in H_2SO_4 as described by (Zhang *et al.* (2008). MDA content was measured by thiobarbituric acid reaction as described by Hodges *et al.* (1999). H_2O_2 content was determined according to Okuda *et al.* (1991).

The whole experiment was repeated five times with 30 seedlings in each repetition. The means were compared by one-way analysis of variance and Duncan's multiple range test at the 5 % level of significance.

After treatment of *Arabidopsis* WT plants with JA for 24 or 48 h, H_2S content in leaves significantly increased

and also the activities of APX, GR, DHAR, MDHAR, GalLDH, and γ -ECS, compared to the control (Tables 1 and 2). Compared with JA alone, application of HT + JA significantly reduced H_2S content and the activities of APX, GR, DHAR, MDHAR, GalLDH and γ -ECS after 24 or 48 h. Exogenous JA had no obvious effects on H_2S content and did not increased activities of the above mentioned enzymes in the H_2S synthesis mutant. Meanwhile, application of HT alone significantly reduced H_2S content and the activities of APX, GR, DHAR, MDHAR, GalLDH and γ -ECS after treatment of 24 and 48 h, compared with control. These results suggest that H_2S may be involved in the regulation of the AsA and GSH metabolism by exogenous JA in the leaves of *A. thaliana*.

Compared to the control, JA significantly increased the ratios of AsA/DHA and GSH/GSSG in the leaves of *Arabidopsis* WT plants after 24 and 48 h (Table 1) but had no obvious effects on these ratios in the H_2S synthesis mutant. Meanwhile, application JA + HT or HT alone significantly reduced the ratio of AsA/DHA and had no obvious effect on the ratio of GSH/GSSG, compared with control. These results suggest that the accumulation of H_2S induced by exogenous JA participated in the regulation of AsA/DHA ratio.

Compared with the control, JA significantly reduced the content of MDA and H_2O_2 in the leaves of *Arabidopsis* WT plants after 24 and 48 h (Table 1). Compared with JA alone, application of HT + JA significantly increased the content of MDA and H_2O_2 . Exogenous JA had no obvious effects on the content of MDA and H_2O_2 in the H_2S synthesis mutant. Meanwhile,

Table 1. Effects of jasmonic acid (JA) and hypotaurine (HT) on the content of H_2S , malondialdehyde (MDA), and H_2O_2 , the ascorbate (AsA)/dehydroascorbate (DHA) and reduced glutathione(GSH)/oxidized glutathione (GSSG) ratios, and activities of cysteine desulfhydrases (*L*-CDes and *D*-CDes) in the leaves of *A. thaliana*. Control - half-strength Hoagland's solution; JA_{WT} - treatment of *Arabidopsis* wild-type with 30 μM JA; HT + JA_{WT} - treatment of *Arabidopsis* wild-type with 20 μM HT + 30 μM JA; JA_{MT} - treatment of H_2S synthetic mutant with 30 μM JA. The plants were pretreated with HT for 8 h, and then exposed to JA for 48 h. Means $\pm$ SEs $n = 5$, different small letters stand for significant differences among different treatments at $P < 0.05$. Differences between treatments lasted 24 and 48 h were not significant.

Parameters	Time [h]	Control	HT	JA _{WT}	HT + JA _{WT}	JA _{MT}
H_2S	24	$6.62 \pm 0.71\text{b}$	$2.15 \pm 0.30\text{c}$	$14.58 \pm 1.35\text{a}$	$7.22 \pm 0.68\text{b}$	$6.59 \pm 0.52\text{b}$
[nmol g ⁻¹ (f.m.)]	48	$7.34 \pm 0.84\text{b}$	$2.44 \pm 0.26\text{c}$	$11.29 \pm 1.11\text{a}$	$6.54 \pm 0.74\text{b}$	$6.90 \pm 0.80\text{b}$
AsA/DHA	24	$18.51 \pm 2.13\text{b}$	$16.87 \pm 1.81\text{c}$	$20.74 \pm 1.75\text{a}$	$17.96 \pm 1.71\text{b}$	$18.06 \pm 1.94\text{b}$
	48	$20.03 \pm 2.01\text{b}$	$18.00 \pm 2.06\text{c}$	$22.95 \pm 2.33\text{a}$	$19.73 \pm 2.51\text{b}$	$19.80 \pm 2.37\text{b}$
GSH/GSSG	24	$20.09 \pm 1.95\text{a}$	$19.81 \pm 2.13\text{a}$	$20.46 \pm 2.03\text{a}$	$19.87 \pm 1.78\text{a}$	$20.30 \pm 2.51\text{a}$
	48	$22.17 \pm 2.46\text{a}$	$21.79 \pm 2.37\text{a}$	$22.55 \pm 2.97\text{a}$	$21.64 \pm 2.16\text{a}$	$22.07 \pm 2.22\text{a}$
MDA	24	$2.19 \pm 0.24\text{b}$	$2.88 \pm 0.35\text{a}$	$1.67 \pm 0.15\text{c}$	$2.11 \pm 0.20\text{b}$	$2.08 \pm 0.23\text{b}$
[nmol g ⁻¹ (f.m.)]	48	$2.41 \pm 0.28\text{b}$	$3.10 \pm 0.33\text{a}$	$1.80 \pm 0.18\text{c}$	$2.32 \pm 0.25\text{b}$	$2.38 \pm 0.21\text{b}$
H_2O_2	24	$6.07 \pm 0.72\text{b}$	$6.97 \pm 0.81\text{a}$	$4.64 \pm 0.51\text{c}$	$5.97 \pm 0.53\text{b}$	$6.12 \pm 0.55\text{b}$
[$\mu\text{mol g}^{-1}$ (f.m.)]	48	$6.53 \pm 0.66\text{b}$	$7.23 \pm 0.71\text{a}$	$4.59 \pm 0.53\text{c}$	$6.35 \pm 0.47\text{ab}$	$6.37 \pm 0.78\text{b}$
<i>L</i> -CDes	24	$13.96 \pm 1.56\text{b}$	$14.83 \pm 1.42\text{b}$	$33.57 \pm 4.11\text{a}$	$35.09 \pm 4.72\text{a}$	$0.00 \pm 0.00\text{c}$
[U mg ⁻¹ (protein)]	48	$15.00 \pm 1.72\text{b}$	$14.15 \pm 1.51\text{b}$	$29.35 \pm 2.75\text{a}$	$30.64 \pm 3.83\text{a}$	$0.00 \pm 0.00\text{c}$
<i>D</i> -CDes	24	$7.25 \pm 1.03\text{b}$	$7.40 \pm 0.97\text{b}$	$16.18 \pm 1.73\text{a}$	$16.43 \pm 1.57\text{a}$	$17.54 \pm 1.77\text{a}$
[U mg ⁻¹ (protein)]	48	$6.81 \pm 0.94\text{b}$	$7.03 \pm 0.85\text{b}$	$18.00 \pm 1.80\text{a}$	$17.55 \pm 1.84\text{a}$	$18.32 \pm 1.95\text{a}$

Table 2. Effects of jasmonic acid (JA) and hypotaurine (HT) on the activities of ascorbate peroxidase (APX), glutathione reductase (GR), dehydroascorbate reductase (DHAR), monodehydroascorbate reductase (MDHAR), γ -glutamylcysteine synthetase (γ -ECS), and L-galactono-1,4-lactone dehydrogenase (GalLDH) in the leaves of *A. thaliana*. Control - half-strength Hoagland's solution; JA_{WT} - treatment of *Arabidopsis* wild-type with 30 μ M JA; HT + JA_{WT} - treatment of *Arabidopsis* wild-type with 20 μ M HT + 30 μ M JA; JA_{MT} - treatment of H₂S synthetic mutant with 30 μ M JA. The plants were pretreated with HT for 8 h, and then exposed to JA for 48 h. Means $\pm$ SEs $n = 5$, different small letters stand for significant differences among different treatments at $P < 0.05$. Differences between treatments lasted 24 and 48 h were not significant.

Parameters	Time [h]	Control	HT	JA _{WT}	HT + JA _{WT}	JA _{MT}
APX	24	0.93 $\pm$ 0.10c	0.45 $\pm$ 0.05d	2.42 $\pm$ 0.26a	1.50 $\pm$ 0.13b	1.02 $\pm$ 0.09c
[U mg ⁻¹ (protein)]	48	0.75 $\pm$ 0.06c	0.37 $\pm$ 0.05d	2.00 $\pm$ 0.22a	1.25 $\pm$ 0.15b	0.81 $\pm$ 0.08c
GR	24	2.44 $\pm$ 0.27b	1.03 $\pm$ 0.15c	3.16 $\pm$ 0.34a	2.30 $\pm$ 0.22b	2.19 $\pm$ 0.24b
[U mg ⁻¹ (protein)]	48	2.25 $\pm$ 0.25b	0.79 $\pm$ 0.09c	3.50 $\pm$ 0.31a	2.25 $\pm$ 0.22b	2.40 $\pm$ 0.21b
DHAR	24	0.62 $\pm$ 0.07b	0.28 $\pm$ 0.04c	1.24 $\pm$ 0.11a	0.71 $\pm$ 0.09b	0.57 $\pm$ 0.05b
[U mg ⁻¹ (protein)]	48	0.50 $\pm$ 0.04b	0.23 $\pm$ 0.03c	1.00 $\pm$ 0.11a	0.60 $\pm$ 0.06b	0.54 $\pm$ 0.04b
MDHAR	24	0.87 $\pm$ 0.09c	0.39 $\pm$ 0.05d	1.60 $\pm$ 0.14a	1.26 $\pm$ 0.12b	1.00 $\pm$ 0.11c
[U mg ⁻¹ (protein)]	48	1.00 $\pm$ 0.11b	0.34 $\pm$ 0.03c	1.65 $\pm$ 0.18a	1.15 $\pm$ 0.11b	1.10 $\pm$ 0.12b
γ -ECS	24	2.88 $\pm$ 0.30c	1.08 $\pm$ 0.15d	5.12 $\pm$ 0.55a	3.84 $\pm$ 0.41b	3.00 $\pm$ 0.25c
[U mg ⁻¹ (protein)]	48	3.00 $\pm$ 0.27c	1.32 $\pm$ 0.21d	5.53 $\pm$ 0.61a	4.22 $\pm$ 0.46b	3.28 $\pm$ 0.35c
GalLDH	24	1.00 $\pm$ 0.12b	0.44 $\pm$ 0.06c	1.93 $\pm$ 0.23a	1.13 $\pm$ 0.13b	1.20 $\pm$ 0.15b
[U mg ⁻¹ (protein)]	48	1.07 $\pm$ 0.12b	0.53 $\pm$ 0.05c	1.70 $\pm$ 0.18a	1.22 $\pm$ 0.11b	1.12 $\pm$ 0.11b

application of HT alone significantly increased the content of MDA and H₂O₂, compared with the control. These results suggested that H₂S has important role in regulating the redox potential in the cells.

In order to elucidate the role of H₂S synthetases L-CDes and D-CDes in regulating the content of H₂S, the effects of different treatments on the activities of L-CDes and D-CDes were studied. The results showed that JA significantly increased the activities of L-CDes and D-CDes in the leaves of *Arabidopsis* WT plants after 24 and 48 h, compared with the control (Table 1). Compared with JA alone, application of HT + JA had no obvious effects on the activities of L-CDes and D-CDes. Meanwhile, application of HT alone had no obvious effects on the activities of L-CDes and D-CDes, compared with control. Exogenous JA also increased D-CDes activity in the H₂S synthesis mutant. However, L-CDes activity was not detected in this mutant. These results suggested that HT has no obvious effect on the activities of H₂S synthetases, and JA could regulate the content of H₂S through regulation of L-CDes and D-CDes.

For the same treatment, there were no significant differences in H₂S content, the activities of L-CDes, D-CDes, APX, GR, DHAR, MDHAR, GalLDH and γ -ECS, the ratios of AsA/DHA and GSH/GSSG, and the content of MDA and H₂O₂ between 24 and 48 h.

H₂S has important roles in improving drought, heat, tolerance and flooding tolerance (Ziogas *et al.* 2015, Chen *et al.* 2016, Li *et al.* 2016, Peng *et al.* 2016). Especially, H₂S could regulate antioxidant responses in plants (Chen *et al.* 2013). For example, Ziogas *et al.* (2015) reported that H₂S could alleviate the damage

induced by drought stress in citrus through antioxidant responses indicated by decreased ion leakage and lipid peroxidation in plants. It has been documented that H₂S could regulate the redox states of AsA and GSH by affecting their biosynthesis and regeneration (Zhang *et al.* 2010b, Shan *et al.* 2011b, Wang *et al.* 2012). Our previous studies (Shan and Liang 2010, Shan *et al.* 2011a) showed that JA also regulated the redox state of AsA and GSH by affecting their biosynthesis and regeneration. It has been shown that endogenous H₂S is involved in the JA-induced stomatal closure (Hou *et al.* 2011). In the present study, we found that exogenous JA increased H₂S content which was involved in the regulation of the AsA and GSH metabolism through increased APX, DHAR, MDHAR, and GalLDH activities which, in turn, increased the ratio of AsA/DHA. Besides, we found that JA-induced H₂S had significant effects on the activities of GR and γ -ECS, but had no obvious effect on the GSH/GSSG ratio. What is the reason for this phenomenon? This may be due to that there exist other regulatory factors, such as other enzymes in the GSH metabolism, including other enzymes in GSH biosynthesis, glutathione peroxidase (GPX) and glutathione S-transferase (GST). These results suggested that H₂S has an important role in JA signalling in regulating the metabolism of AsA and GSH and the ratio of AsA/DHA in plants, which was the new finding in comparison with the already published data.

L-CDes and D-CDes are two enzymes known to produce H₂S in plants (Yamasaki and Cohen 2016). It was not surprising that the *Alt-cdes* mutant had no L-CDes activity, but it had higher D-CDes activity in response to JA, compared with control. Meanwhile, JA

also significantly increased the activities of *L*-CDes and *D*-CDes in WT plants. Thus, we have strong evidence that the activities of *L*-CDes and *D*-CDes were induced by JA among the other enzymes that produce H₂S in *Arabidopsis*, which may be also a new finding.

As a signal molecule, JA has important role in modulating the redox state of the cells through the AsA and GSH metabolism. However, the relationship between H₂S, NO, and MEK1/2 in the regulation of the AsA and

GSH metabolism during JA signalling remains unclear. Therefore, it might be very interesting to investigate the relationships between JA, H₂S, H₂O₂, NO, and MEK1/2 in the regulation of the AsA and GSH metabolism in detail.

In conclusion, the results clearly suggest that exogenous JA-induced H₂S accumulation participated in the regulation of the AsA and GSH metabolism, which, in turn, maintains the redox state in the cells.

References

- Bradford, M.M.: A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. - *Anal. Biochem.* **72**: 248-254, 1976.
- Chen, J., Shang, Y.T., Wang, W.H., Chen, X.Y., He, E.M., Zheng, H.L., Shanguan Z.: Hydrogen sulfide-mediated polyamines and sugar changes are involved in hydrogen sulfide-induced drought tolerance in *Spinacia oleracea* seedlings. - *Front. Plant Sci.* **7**: 1-18, 2016.
- Chen, J., Wang, W.H., Wu, F.H., You, C.Y., Liu, T.W., Dong, X.J., He, J.X., Zheng, H.L.: Hydrogen sulfide alleviates aluminum toxicity in barley seedlings. - *Plant Soil* **362**: 301-318, 2013.
- Dalton, D.A., Russell, S.A., Hanus, F.J., Pascoe, G.A., Evans, H.J.: Enzymatic reactions of ascorbate and glutathione that prevent peroxide damage in soybean root nodules. - *Proc. nat. Acad. Sci. USA* **83**: 3811-3815, 1986.
- Grace, S.C., Logan, B.A.: Acclimation of foliar antioxidant systems to growth irradiance in three broad-leaved evergreen species. - *Plant Physiol.* **112**: 1631-1640, 1996.
- Griffith, O.W.: Determination of glutathione and glutathione disulfide using glutathione reductase and 2-vinylpyridine. - *Anal. Biochem.* **106**: 207-212, 1980.
- Hodges, M.D., DeLong, J.M., Forney, C.F., Prange, R.K.: Improving the thiobarbituric acid-reactive-substances assay for estimating lipid peroxidation in plant tissues containing anthocyanin and other interfering compounds. - *Planta* **207**: 604-611, 1999.
- Hodges, D.M., Andrews, C.J., Johnson, D.A., Hamilton, R.I.: Antioxidant compound responses to chilling stress in differentially sensitive inbred maize lines. - *Physiol. Plant.* **98**: 685-692, 1996.
- Hou, Z.H., Liu, J., Hou, L.X., Li, X.D., Liu, X.: H₂S may function downstream of H₂O₂ in jasmonic acid-induced stomatal closure in *Vicia faba*. - *Chin. Bull. Bot.* **46**: 396-406, 2011.
- Li, L., Wang, Y., Shen, W.: Roles of hydrogen sulfide and nitric oxide in the alleviation of cadmium-induced oxidative damage in alfalfa seedling roots. - *Biometals* **25**: 617-631, 2012.
- Li, Z.G., Gu, S.P.: Hydrogen sulfide as a signal molecule in hematin-induced heat tolerance of tobacco cell suspension. - *Biol. Plant.* **60**: 595-600, 2016.
- Liu, J., Hou, L.X., Liu, G.H., Liu, X., Wang, X.C.: Hydrogen sulfide induced by nitric oxide mediates ethylene-induced stomatal closure of *Arabidopsis thaliana*. - *Chin. Sci. Bull.* **56**: 2515-2522, 2011.
- Miyake, C., Asada, K.: Thylakoid-bound ascorbate peroxidase in spinach chloroplasts and photoreduction of its primary oxidation product monodehydroascorbate radicals in thylakoids. - *Plant Cell Physiol.* **33**: 541-553, 1992.
- Nakano, Y., Asada, K.: Hydrogen peroxide is scavenged by ascorbate specific peroxidase in spinach chloroplasts. - *Plant Cell Physiol.* **22**: 867-880, 1981.
- Noctor, G., Foyer, C.H.: Ascorbate and glutathione: keeping active oxygen under control. - *Annu. Rev. Plant Physiol.* **49**: 249-279, 1998.
- Okuda, T., Masuda, Y., Yamanka, A., Sagisaka, S.: Abrupt increase in the level of hydrogen peroxide in leaves of winter wheat is caused by cold treatment. - *Plant Physiol.* **97**: 1265-1267, 1991.
- Peng, R., Bian, Z., Zhou, L., Cheng, W., Hai, N., Yang, C., Yang, T., Wang, X., Wang, C.: Hydrogen sulfide enhances nitric oxide-induced tolerance of hypoxia in maize (*Zea mays* L.). - *Plant Cell Rep.* **35**: 2325-2340, 2016.
- Radhika, V., Kost, C., Boland, W., Heil, M.: The Role of jasmonates in floral nectar secretion. - *PloS ONE* **5**: 9265, 2010.
- Ren, C., Dai, C.: Jasmonic acid is involved in the signaling pathway for fungal endophyte-induced volatile oil accumulation of *Atractylodes lancea* plantlets. - *BMC Plant Biol.* **12**: 128-133, 2012.
- Riemenschneider, A., Nikiforova, V., Hoefgen, R., Papenbrock, J.: Impact of elevated H₂S on metabolite levels, activity of enzymes and expression of genes involved in cysteine metabolism. - *Plant Physiol. Biochem.* **43**: 473-483, 2005.
- Rüeggsegger, A., Brunold, C.: Effect of cadmium on γ -glutamylcysteine synthesis in maize seedlings. - *Plant Physiol.* **99**: 428-433, 1992.
- Shan, C., Liang, Z.: Jasmonic acid regulates ascorbate and glutathione metabolism in *Agropyron cristatum* leaves under water stress. - *Plant Sci.* **178**: 130-139, 2010.
- Shan, C., Liang, Z., Sun, Y., Hao, W., Han, R.: The protein kinase MEK1/2 participates in the regulation of ascorbate and glutathione content by jasmonic acid in *Agropyron cristatum* leaves. - *J. Plant Physiol.* **168**: 514-518, 2011a.
- Shan, C., Zhang, S., Li, D., Zhao, Y., Tian, X., Zhao, X., Wu, Y., Wei, X., Liu, R.: Effects of exogenous hydrogen sulfide on the ascorbate and glutathione metabolism in wheat seedlings leaves under water stress. - *Acta Physiol. Plant.* **33**: 2533-2540, 2011b.
- Shan, C., Zhou, Y., Liu, M.: Nitric oxide participates in the regulation of the ascorbate-glutathione cycle by exogenous jasmonic acid in the leaves of wheat seedlings under drought stress. - *Protoplasma* **252**: 1397-1405, 2015.
- Wang, L., Hou, Z., Hou, L., Zhao, F., Liu, X.: H₂S induced by

- H₂O₂ mediates drought-induced stomatal closure in *Arabidopsis thaliana*. - Chin. Bull. Bot. **47**: 217-225, 2012.
- Yamasaki, H., Cohen, M.F.: Biological consilience of hydrogen sulfide and nitric oxide in plants: gases of primordial earth linking plant, microbial and animal physiologies. - Nitric Oxide **55-56**: 91-100, 2016.
- Zhang, H., Hu, L.Y., Hu, K.D., He, Y.D., Wang, S.H., Luo, J.P.: Hydrogen sulfide promotes wheat seed germination and alleviates the oxidative damage against copper stress. - J. Integr. Plant Biol. **50**: 1518-1529, 2008.
- Zhang, H., Hu, L.Y., Li, P., Hu, K.D., Jiang, C.X., Luo, J.P.: Hydrogen sulfide alleviated chromium toxicity in wheat. - Biol. Plant. **54**: 743-747, 2010b.
- Zhang, H., Tang, J., Liu, X.P., Wang, Y., Yu, W., Peng, W.Y., Fang, F., Ma, D.F., Wei, Z.J., Hu, L.Y.: Hydrogen sulfide promotes root organogenesis in *Ipomoea batatas*, *Salix matsudana* and *Glycine max*. - J. Integr. Plant Biol. **51**: 1086-1094, 2009a.
- Zhang, H., Wang, M.J., Hu, L.Y., Wang, S.H., Hu, K.D., Bao, L.J., Luo, J.P.: Hydrogen sulfide promotes wheat seed germination under osmotic stress. - Russ. J. Plant Physiol. **57**: 532-539, 2010a.
- Zhang, H., Ye, Y.K., Wang, S.H., Luo, J.P., Tang, J., Ma, D.F.: Hydrogen sulfide counteracts chlorophyll loss in sweetpotato seedling leaves and alleviates oxidative damage against osmotic stress. - Plant Growth Regul. **58**: 243-250, 2009b.
- Ziogas, V., Tanou, G., Belghazi, M., Filippou, P., Fotopoulos, V., Grigorios, D., Molassiotis, A.: Roles of sodium hydrosulfide and sodium nitroprusside as priming molecules during drought acclimation in citrus plants. - Plant mol. Biol. **89**: 433-450, 2015.